

537,963

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
22 July 2004 (22.07.2004)

PCT

(10) International Publication Number  
**WO 2004/060052 A2**

- (51) International Patent Classification<sup>7</sup>: **A01K**
- (21) International Application Number:  
PCT/US2003/039686
- (22) International Filing Date:  
11 December 2003 (11.12.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/434,115 16 December 2002 (16.12.2002) US  
60/476,481 5 June 2003 (05.06.2003) US
- (71) Applicant (for all designated States except US): **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **CHAN, Andrew, Chee-Yuen** [US/US]; 1201, Cloud Avenue, Menlo Park, CA 94025 (US). **GONG, Qian** [CN/US]; 755 Neptune Lane, Foster City, CA 94404 (US). **MARTIN, Flavius** [RO/US]; 730 Bounty Dr., #3004, Foster City, CA 94404 (US).
- (74) Agent: **BRUESS, Steven, C.**; Merchant & Gould P.C., P.O. Box 2903, Minneapolis, MN 55402-0903 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TRANSGENIC MICE EXPRESSING HUMAN CD20 AND/OR CD16

(57) Abstract: The present invention generally relates to non-human transgenic animals expressing human cellular markers, including CD20 and/or preferably CD16.

WO 2004/060052 A2

BEST AVAILABLE COPY

## TRANSGENIC MICE EXPRESSING HUMAN CD20 AND/OR CD16

This application is being filed as a PCT International Patent Application in the name of Genentech, Inc., a U.S. national corporation and resident, (Applicant for all countries except US); Andrew Chee-Yuen Chan, a U.S. citizen and resident (Applicant for US only); Qian Gong, a Chinese citizen and U.S. resident (Applicant for US only); and Flavius Martin, a Romanian citizen and U.S. resident (Applicant for US only), on 11 December 2003, designating all countries and claiming priority to United States provisional application Serial Number 60/434,115, filed December 16, 2002, and United States provisional application Serial Number 60/476,481, filed June 5, 2003.

### Background of the Invention

T and B cells both comprise cell surface proteins that can be utilized as markers for differentiation and identification. One such human B cell marker is the human B lymphocyte-restricted differentiation antigen Bp35, also known as "CD20". CD20 is expressed during early pre-B cell development and remains until plasma cell differentiation. It is believed that the CD20 molecule regulates a step in the activation process that is required for cell cycle initiation and differentiation and is usually expressed at very high levels on neoplastic B cells.

CD20 is present on both normal B cells as well as malignant B cells, whose unabated proliferation can lead to B cell lymphoma. Thus, the CD20 surface antigen has the potential of serving as a candidate for targeting of B cell lymphomas with antibodies specific to the antigen. These anti-CD20 antibodies specifically bind to the CD20 cell surface antigen of both normal and malignant B cells, leading to the destruction and depletion of B cells. Chemical agents or radioactive labels having the potential to destroy the tumor can be conjugated to the anti-CD20 antibody such that the agent is specifically delivered to the neoplastic B cell.

The use of monoclonal antibodies targeting CD20 has been described (see, for example, Weiner, Semin. Oncol., 26, 43-51 (1999); Gopal and Press, J. Lab. Clin. Med., 134, 445-450 (1999); White et al., Pharm. Sci. Technol. Today, 2, 95-101 (1999)). Rituxan<sup>TM</sup> is a chimeric anti-CD20 monoclonal antibody that has been used widely both as a single agent and together with chemotherapy in patients with newly diagnosed and relapsed lymphomas (Davis et al, J. Clin. Oncol., 17, 1851-

1857 (1999); Solal-Celigny et al., *Blood*, 94, abstract 2802 (1999); Foran et al., *J. Clin. Oncol.*, 18, 317-324 (2000). The use of radiolabeled antibody conjugates has also been described (for example, Bexxar<sup>TM</sup>; Zelenetz et al., *Blood*, 94, abstract 2806 (1999)).

5           The interaction of antibody-antigen complex with cells of the immune system results in a wide array of responses, ranging from effector functions such as antibody-dependent cytotoxicity, mast cell degranulation, and phagocytosis to immunomodulatory signals such as regulating lymphocyte proliferation and antibody secretion. All these interactions are initiated through the binding of the Fc  
10 domain of antibodies or immune complexes to specialized cell surface receptors on hematopoietic cells. It is now well established that the diversity of cellular responses triggered by antibodies and immune complexes results from the structural heterogeneity of Fc receptors (FcRs).

One group of these receptors, FcγRs, is found on most cells of the  
15 hematopoietic lineage, and mediate both high and low affinity binding to IgG (see, for example, U.S. Patent No. 5,877,396, incorporated herein by reference). The high affinity receptor, FcγRI, binds monomeric IgG and is expressed exclusively on macrophages and neutrophils. It is capable of mediating antibody-dependent cell-mediated cytotoxicity (ADCC) and phagocytosis in response to crosslinking by  
20 antibody. The low affinity receptors for IgG, FcγRII and FcγRIII (CD16), are responsible for effector cell responses to immune complexes and represent the FcγRs primarily involved in the inflammatory response in vivo. FcγRII is widely expressed on haematopoietic cells and functions as an inhibitory receptor on B cells, while on cells of the myeloid lineage and on platelets, FcγRII triggers ADCC,  
25 phagocytosis and the release of inflammatory mediators when crosslinked by immune complexes. FcγRIII is expressed on various leucocytes including natural killer (NK) cells, macrophages, neutrophils, eosinophils, basophils and mast cells, and mediates effector responses when crosslinked by immune complexes. It is the sole FcR on NK cells, mediating all the antibody-dependent responses on those  
30 cells. Natural killer cells are a subset of spontaneously cytotoxic lymphocytes that lytically destroy tumor cells without apparent antigen specificity or restriction by histocompatibility molecules. In addition to these well-characterized effector cell pathways, FcγRIII has been found on immature thymocytes, where it has been postulated to function in early thymocyte development.

With other receptors of the immunoglobulin Fc portion (eg. FcγRI, FcγRII, FcεRI), CD16 plays an important role in mediating autoimmunity and inflammatory responses. Studies using monoclonal antibodies against CD16 have established this receptor's role in removing immune complexes from circulation and in mediating

5 ADCC (see for example Van de Winkel et al., Immunol. Today, 14, 215-221 (1993)). The binding of IgG with CD16 elicits NK/LGL cell activation and triggers ADCC. ADCC can be halted in the presence of high levels of soluble CD16.

It has been found (see Mathiot et al., J. Clin. Immunol., 13, 41-8 (1993)) that the level of soluble CD16 was significantly decreased in patients with multiple

10 myeloma compared with healthy volunteers. In addition a stage-dependent decrease of soluble CD16 was observed, with a highly significant difference between stage I and stages II+III myeloma patients. Therefore, measurement of soluble CD16 in serum is both a diagnostic and a prognostic marker of myeloma, which can be useful to define and guide novel immunomodulatory therapies of the disease.

It has further been found that CD16 is present in human serum and other body fluids and is elevated at sites of inflammation (see Fleit et al., Blood, 79, 2721-8 (1992)). It appears that there are at least two forms of human CD16, type A and type B. CD16-A is expressed predominantly on the surface of macrophages, natural killer cells and large granular lymphocytes (NK/LGL), whereas CD16-B is

15 20 expressed predominantly on the surface of neutrophils and monocytes.

In spite of the significant roles of CD20 and CD16 in human lymphoma and in inducement of important immunological responses, animal models are lacking which co-express the human markers. Thus, a need exists for relevant animal models for disease study and pharmaceutical drug development.

25

### Summary of the Invention

The present invention generally relates to non-naturally occurring non-human transgenic animals expressing human cellular markers, specifically, CD16 and CD20. In one aspect, the transgenic animals provide a system to identify and

30 test novel therapeutic agents for CD20 associated disease or conditions, such as cancer. In an embodiment, the transgenic animals are useful to test efficacy and toxicities of CD20 directed therapies.

The invention provides a non-naturally occurring transgenic animal whose genome comprises a nucleotide sequence encoding a heterologous CD20, preferably,

human CD20. The nucleotide sequence is preferably operably linked to a human endogenous promoter whereby the human CD20 is expressed on the surface of B lymphocytes. In one embodiment, the human CD20 transgenic mice are characterized by expression of human CD20 on cells at a level sufficient for anti-  
5 human CD20 antibody bound to the expressing cells to affect killing of the cells, resulting in B cell depletion of peripheral and/or circulating B cells of at least about 75% and more preferably, 80%, 85%, 90%, 95% , 99% and even 100%.

In one embodiment of the invention, the genome of the non- naturally occurring non-human transgenic animal further comprises a nucleotide sequence  
10 encoding a heterologous FcγIII receptor, preferably, human CD16 and preferably, the α chain of human CD16. The nucleotide sequence is preferably operably linked to a human endogenous promoter, whereby the heterologous receptor is expressed on the surface of leukocytes, including one or more of the following: natural killer (NK) cells, macrophages, neutrophils, eosinophils, basophils, thymocytes, and mast  
15 cells.

According to a preferred embodiment, when the genome of the animal comprises a homologous endogenous gene (either CD20 or CD16, or both), the gene is disrupted or knocked out such that endogenous molecule is not expressed on cell surfaces.

20 The present invention further provides methods of identifying agents capable of treating B cell lymphoma where the method comprises administering an agent to a transgenic animal that expresses human CD20 on B lymphocytes and determining whether there is a reduction in the number of B lymphocytes. The invention also provides methods of identifying agents capable of depleting or killing cells  
25 expressing human CD20 comprising administering an agent to transgenic animals that express human CD20 and determining whether there is a reduction in the number of such cells. Further provided are agents identified according to such methods.

The animal models of the present invention may also be used to identify  
30 agents capable of inducing effector cell responses such as ADCC or NK cell mediated immune responses. After administration of the agent, the animal can be monitored for an immune response such as by determining increase or decrease in cytokine levels. An increase in levels of cytokines after administration of the agent identifies an agent that induces a Fc- mediated effector cell response. Binding of the

putative agent to CD16 can also be assessed using suitable markers or labels. Additionally, agents can be screened for their ability to affect depletion of B cells expressing human CD20 (including malignant cells) via CD16 mediated immune response by comparing B cell depletion in the CD20<sup>+</sup> transgenic animal with that in  
5 the CD16<sup>+</sup>CD20<sup>+</sup> transgenic animal after administration of an agent. The animals of the present invention are also useful for assessing the toxicity of anti-CD20 therapeutics by administration to the presently described transgenic animals.

Treatment specificity, toxicity and efficacy can also be determined by comparison of the agent's effect with that in a wild-type animal or untreated  
10 transgenic animal. A non-human transgenic animal of the present invention can further provide an indication of the safety of a particular agent for administration to a human. For example, a humanized antibody or other agent can be administered to the transgenic animal and any toxic or adverse effects as a result of the administration of the agent to the animal can be monitored as an indication of the  
15 safety and tolerability of the humanized antibody or agent for in vivo human use. Adverse events that may occur on a short term basis include headache, infection, fever, chills, pain, nausea, asthenia, pharyngitis, diarrhea, rhinitis, infusion reactions, and myalgia. Short term adverse events are measured in days post treatment. Long  
20 term adverse effects include cytotoxicity of certain cell types, bleeding events due to thrombocytopenia, release of mediators due to inflammatory and/or allergic reactions, inhibition of the immune system and/or development of an anti-therapeutic agent antibody, end organ toxicity, and increased incidence of infection or malignancy. Long term adverse events are measured in months post treatment.

Another aspect of the invention involves a method for determining efficacy  
25 of an anti-CD20 agent. Efficacy can be determined by administering a range of doses of the agent to set of transgenic animals having human CD20 and/or human CD16 alpha chain, determining at least one dose of the agent that results in a decrease in cells bearing human CD20.

### 30 **Brief Description of the Figures**

Figure 1 shows the expression of human CD20 in mouse B220<sup>+</sup> cells derived from mice without the transgene (Tg-), heterozygous (Tg+/-) and homozygous (Tg+/+) for the transgene.

Figure 2 provides a schematic diagram of expression of various cell surface markers (CD43, IgM, IgD) during B cell differentiation and maturation. In the Tg<sup>+</sup> mice, hCD20 is expressed on pre-B, immature B cells and mature B cells.

Figure 3 shows the result of screening Tg<sup>+</sup> mice for human CD20 expression in the B cells of the bone marrow. Cells were stained with anti-human CD20 conjugated to FITC (BD Pharmingen). Gating the cells for presence of B220 and CD43 allows delineation into the various populations of B cells. For gating, cells were stained with anti-B220 Ab conjugated to PerCP (BD Pharmingen) and with anti-CD43 Ab conjugated to PE (fluorescence, BD Pharmingen).

Figure 4 shows the result of screening Tg<sup>+</sup> mice for human CD20 expression in the B cells of the spleen. Cells were stained with anti-human CD20 conjugated to FITC (BD Pharmingen). Gating the cells for B220 and CD21 allows delineation into the various populations of B cells. For gating, cells were stained with anti-B220 Ab conjugated to PerCP (BD Pharmingen) and with anti-CD21 Ab conjugated to PE (fluorescence, BD Pharmingen). B cells with human CD20 are found in the T1 zone, marginal zone and T2/Follicular zone.

Figure 5 shows the result of screening Tg<sup>+</sup> mice for human CD20 expression in the B cells of the mesenteric lymph nodes. Cells were stained with anti-human CD20 conjugated to FITC (BD Pharmingen). Gating the cells for B220 and CD21 allows delineation into the various populations of B cells. For gating, cells were stained with anti-B220 Ab conjugated to PerCP (BD Pharmingen) and with anti-CD21 Ab conjugated to PE (fluorescence, BD Pharmingen).

Figure 6 shows the result of screening Tg<sup>+</sup> mice for human CD20 expression in the B cells of the Peyer's Patches. Cells were stained with anti-human CD20 conjugated to FITC (BD Pharmingen). Gating the cells for B220 and CD38 allows delineation into the various populations of B cells. For gating, cells were stained with anti-B220 Ab conjugated to PerCP (BD Pharmingen) and with anti-CD38 Ab conjugated to PE (fluorescence, BD Pharmingen). B cells with human CD20 are mature B cells and cells in the germinal centers.

Figure 7 is a schematic diagram of administration of anti-human CD20 mAb into Tg<sup>+</sup> mice and analysis of presence or absence of cells having human CD20. A single dose of 1 mg of anti-human CD20 monoclonal antibody was administered at day 0. Samples were taken from various tissues at day -1, day 1, 2, 3, 4, 7, 14 and 21. Samples from different tissues such as peripheral blood spleen, lymph nodes,

bone marrow, and Peyer's Patches were analyzed by FACs as described previously. Serum levels of anti-CD20 monoclonal antibody were also monitored.

Figure 8 shows depletion of peripheral B cells in transgenic mice treated with anti-human CD20 mAb m2H7 (BD Pharmingen). The antibodies were  
5 administered to the transgenic mice as outlined in the schematic in Figure 7 at a dose of 1 mg total. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches with gating as described previously.

Figure 9 shows depletion of mature peripheral lymph node B cells in transgenic mice treated with anti-human CD20 mAb m2H7. The antibodies were  
10 administered to the transgenic mice as outlined in the schematic in Figure 7 at a dose of 1 mg total. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches with gating as described previously.

Figure 10 shows depletion of splenic T2 and follicular B cells in transgenic mice treated with anti-human CD20 mAb m2H7. The antibodies were administered  
15 to the transgenic mice as outlined in the schematic in Figure 7 at a dose of 1 mg total. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches with gating as described previously.

Figure 11 shows depletion of re-circulating mature B cells in transgenic mice treated with anti-human CD20 mAb m2H7. The antibodies were administered to the  
20 transgenic mice as outlined in the schematic in Figure 7 at a dose of 1 mg total. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches with gating as described previously.

Figure 12 shows depletion of mature B cells and resistance of Peyer's Patches germinal center B cells to depletion in transgenic mice treated with anti-  
25 human CD20 mAb m2H7. The antibodies were administered to the transgenic mice as outlined in the schematic in Figure 7 at a dose of 1 mg total. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches with gating as described previously.

Figure 13 shows depletion and recovery of B cells following administration  
30 of anti-CD20 mAb. Antibody was administered to the mice at day 1. At day 6 post antibody treatment, B cells expressing human CD20 in the peripheral blood were not detectable. At week 6, upon clearance of the antibody, hCD20+ cells begin to be detected. By week 14, B cells appeared to have recovered to normal levels.



Recovery results from precursor B cells which do not express CD20 and that then subsequently develop into mature B cells with human CD20+.

Figure 14 shows FACS plot indicating resistance of splenic germinal center B cells to short-term anti-CD20 mAb therapy. Mice were unimmunized or  
5 immunized with sheep red blood cells (SRBC) by intraperitoneal injection at day 1 to induce germinal centers in the spleen. The germinal centers appear by day 7. At day 8, one group of mice was treated with anti-CD20 antibody m2H7. The control set of mice was treated with mIgG2a isotype control antibody. Spleen cells from the mice were analyzed at day 12. PNA (peanut agglutinin), which stains for germinal  
10 center, was utilized. No detectable germinal center cells were seen in the spleens of mice not immunized with SRBC whereas the spleens of immunized mice show 0.3% PNA staining cells. While T2/Follicular B cells are depleted with anti-CD20 antibody treatment, marginal center B cells in the spleen are resistant to the antibody.

Figure 15 shows a T cell independent response in anti-CD20 mAb treated mice and controls. Mice were treated with m2H7 or isotype control antibody mIgG2a at day 0. At days 3-7, B cell depletion has occurred. At day 7, the mice were injected i.v. with Streptococcus Pneumoniae IV to induce a response to the polysaccharide. A T cell independent response was mounted on day 11. The results  
20 demonstrate that treatment with anti-human CD20 m2H7 did not affect the B cell response from the marginal zone and B 1 cells, i.e., the non-depleted MZ and B1 B cells confer protection to T-independent antigens.

Figure 16 shows a comparison of the effect of BR3 and 2H7 antibody treatment in Tg<sup>+</sup> mice. Human CD20 transgenic mice expressing a bacterial artificial  
25 chromosome encoding human CD20 (designated as hCD20<sup>+</sup> mice) were treated with intraperitoneal injections of anti-CD20 mAb (single injection of 100 micrograms on day 9), BR3-Fc (100 micrograms every other day from days 1 through 12), or the combination of anti-CD20 mAb and BR3-Fc. Each group consisted of 4 mice. Two days following the last injection, the mice were sacrificed and analyzed for hCD20<sup>+</sup>  
30 B cells. FACS analysis of spleen, blood, lymph node and Peyer's Patches were analyzed for B cell markers (CD21<sup>+</sup>CD23<sup>+</sup>). Anti-CD20 mAb therapy depleted T2 and follicular B cells, but not marginal zone B cells in the spleen, whereas BR3-Fc treatment decreased T2/follicular and marginal zone B cells in the spleen.

Figure 17 shows the lack of effect of anti-CD20 mAb treatment on Peyer's Patches. Human CD20 transgenic mice expressing a bacterial artificial chromosome encoding human CD20 (designated as hCD20<sup>+</sup> mice) were treated with intraperitoneal injections of anti-CD20 mAb (single injection of 100 micrograms on day 9), BR3-Fc (100 micrograms every other day from days 1 through 12), or the combination of anti-CD20 mAb and BR3-Fc. Each group consisted of 4 mice. Two days following the last injection, the mice were sacrificed and analyzed for hCD20<sup>+</sup> B cells. FACS analysis of spleen, blood, lymph node and Peyer's Patches were analyzed for B cell markers (CD21<sup>+</sup>CD23<sup>+</sup>). Neither BR3-Fc nor 2H7 nor the combination of the two had an effect on germinal center B cells in Peyer's Patches

Figure 18 shows that plasma cells are not depleted following long-term anti-CD20 mAb treatment. Transgenic Tg<sup>+</sup> mice positive for human CD20 were treated with anti-human CD20 mH27 antibody as described previously. The mice were analyzed for presence or absence of plasma cells by detecting cells in bone marrow and spleen that were positive for syndican (CD-138 plasma cell marker). The number of IgA or IgM positive plasma cells was also monitored after anti-human CD20 treatment. The results show that plasma cells in Tg<sup>+</sup> mice were not affected by the anti-human CD20 antibody treatment, indicating that the Tg<sup>+</sup> mice still have the capacity to generate antibodies.

Figure 19 shows depletion of NK cells by PK-136 mAb in Tg<sup>+</sup> mice. Hybridoma clone, which produces PK-136 mAb (specific against mouse NK1.1), was obtained from ATCC. Four groups of human CD20 transgenic mice were injected ip with control mAb, PK-136, anti-CD20 mAb and the combination of PK-136/anti-CD20, respectively. Doses administered ip were as follows:

control mAb:	200ug/ip, 3 ip/week, for 1 week
PK-136:	200ug/ip, 3 ip/week, for 1 week
anti-CD20 mAb:	10 ug/ip, single dose

NK cells from peripheral blood, liver and spleen were analyzed after treatment with antibodies. Data is expressed as mean $\pm$ standard error, with n=8.

Figure 20 shows that NK cells play a role in 2H7 mediated B cell depletion in Tg<sup>+</sup> mice. Hybridoma clone, which produces PK-136 mAb (specific against mouse NK1.1), was obtained from ATCC. Four groups of human CD20 transgenic mice were injected ip with control mAb, PK-136, anti-CD20 mAb and the

combination of PK-136/anti-CD20, respectively. Doses administered ip were as follows:

- control mAb: 200ug/ip, 3 ip/week, for 1 week  
PK-136: 200ug/ip, 3 ip/week, for 1 week  
5 anti-CD20 mAb: 10 ug/ip, single dose

Lymphocytes from peripheral blood, lymph nodes and spleen were analyzed 3 days after anti-CD20 mAb ip. Data is expressed as mean+/-standard error, with n=8.

Figure 21 shows the expression of human CD20 and human CD16 on different populations of cells in transgenic mice. Blood cells from CD20Tg-  
10 /Cd16Tg- (control mice), CD20Tg+/CD16Tg-, CD20Tg-/CD16Tg+, and CD20Tg+/CD16+ mice were stained with labeled anti-human CD20 antibodies labeled with FITC, anti-B220 antibody conjugated to PerCP and anti-human CD16 antibodies labeled with PE (BD Pharmingen) and analyzed by FACS. The results show that  
15 human CD20 is found on B cells and the human CD16 is found on cells that lack the B220 marker and therefore, are not B cells. Transgenic mice positive for both markers show both populations of cells.

Figure 22A shows a representative amino acid sequence (SEQ ID NO: 1) (GenBank Accession No. NM000569), cDNA for human CD16 alpha chain isotype A (SEQ ID NO: 2) (GenBank Accession No. NM000569) (Figure 22B) and a  
20 genomic DNA sequence for human CD16 alpha chain isotype A (SEQ ID NO: 3) (GenBank Accession No. Z46222) (Figure 22C). Figure 22D shows a representative cDNA sequence for mouse CD16 alpha chain (Gen Bank Accession No. NM010188) (SEQ ID NO: 9). Figure 22E shows a representative amino acid sequence (SEQ ID NO: 10) and cDNA sequence for human CD16  $\alpha$  chain isotype B  
25 (SEQ ID NO: 11) (GenBank Accession No. NM000570). Figure 22F shows a representative genomic sequence encoding human CD16  $\alpha$  chain isotype B (SEQ ID NO: 12) (GenBank Accession No. Z46223). Figure 22G shows a cDNA sequence encoding murine CD16 alpha chain (SEQ ID NO: 13) (GenBank Accession No. NM010188).

30 Figure 23A shows an amino acid sequence for human CD20 (SEQ ID NO: 4) (GenBank Accession No. BC002807), cDNA sequence for human CD20 (Figure 23B) (SEQ ID NO: 5) (GenBank Accession No. BC002807) and a genomic sequence for human CD20 (SEQ ID NO: 6) (GenBank Accession No. AH005353)

(Figure 23C). Figure 23D shows a representative cDNA sequence for murine CD20 (SEQ. ID NO:14) (GenBank Accession No. M62541).

Figure 24 shows a comparison of expression of human CD16 in human CD16Tg<sup>-</sup> mice to human CD16Tg<sup>+</sup> mice. The cells were stained with anti-human  
5 CD16 conjugated to PE (BD Pharmingen) and were also stained with anti-DX5-FITC (BD Pharmingen) to identify NK cells or anti-F4/80 conjugated to APC (Allophycocyanin) to identify macrophages. The results show that both NK and macrophages express the human CD16 transgene in human CD16Tg<sup>+</sup> mice.

Figure 25 shows an amino acid (GenBank Accession No. P30273) (SEQ ID  
10 NO: 7) and cDNA sequence (GenBank Accession No. M33915) (SEQ ID NO: 8) for human Fc receptor gamma chain.

Figure 26 shows an FACS analysis for the presence or absence of expression mouse CD16 on macrophages. Peripheral blood cells from mice lacking murine CD16 (CD16<sup>-/-</sup>) and from control mice having murine CD16 (CD16<sup>+/+</sup>) were stained  
15 with anti-mouse CD16 antibody. The cells were gated for expression of a macrophage marker mac-1 using anti-mac 1 antibodies.

Figure 27 shows a FACS analysis for the presence or absence of expression of mouse CD64 on peripheral blood cells from CD16<sup>-/-</sup> mice. CD64 is the Fcgamma RI and expression of this receptor on mouse cells shows that expression of  
20 other Fc receptors was not affected by knocking out the Fcgamma RIII (CD16) alpha chain. Peripheral blood cells from mice lacking murine CD16 (CD16<sup>-/-</sup>) and from isotype control mice were stained with anti-mouse CD64 antibody. The cells were gated for expression of a macrophage marker mac 1 using anti-mac 1 antibodies.

Figure 28 shows a representative comparison of the expression level of  
25 human CD20 expression on peripheral blood cells from a human CD20 transgenic mouse as compared to expression of CD20 on human peripheral blood cells. Peripheral blood cells were obtained from a human donor and from a hCD20Tg<sup>+</sup> mouse and stained with labeled anti-human CD20 antibody (mH27). The cells were  
30 analyzed by FACS and were gated on human CD19<sup>+</sup> and B220<sup>+</sup> populations. The numbers on the graph represent mean fluorescence intensity.

### Detailed Description

The following terms have the meanings ascribed to them below unless specified otherwise.

The term “construct” or “targeting construct” refers to a polynucleotide molecule that comprises a targeting region. A targeting region comprises a sequence that is substantially homologous to an endogenous sequence in a target tissue, cell or animal and that provides for integration of the targeting construct into the genome of the target tissue, cell or animal. Typically, the targeting construct will also include a gene or a nucleic acid sequence of particular interest, a marker gene and appropriate control sequences.

The term “CD16” or “FcγRIII” are used interchangeably and refer to a cell surface receptor protein for a Fc portion of an IgG immunoglobulin. This receptor is a low affinity receptor for IgG and preferentially binds IgG in immune complexes. FcγRIII receptor is comprised of an  $\alpha$  chain which serves as a ligand binding chain and a homodimer or heterodimer. When the FcγRIII is expressed on macrophages, the  $\alpha$  chain is associated with a homodimer of the gamma chain. When the FcγRIII is expressed on natural killer cells, the  $\alpha$  chain is associated with a heterodimer of the gamma chain with a delta chain. The gamma chain is involved in cell surface expression of the FcγRIII. “Naturally occurring CD16” has the amino acid sequence of cell surface receptor protein obtained from nature and includes naturally occurring variant forms including allelic variants, isotypes and truncated forms. Human CD16 includes all isotypes of the  $\alpha$  chain including both CD16 or FcγRIII-A (GenBank Accession no. Z46222) and CD16 or FcγRIII-B (GenBank Accession no. Z46223). Representative amino acid and nucleotide sequences of the alpha chain of FcγRIII are shown in Figure 22A/B/C/D/E. Representative sequences for the human gamma chain are shown in Figure 25 (GenBank Accession No. P30273 and M33195). The invention also contemplates CD16 or FcγRIII variants. The variants are changed as compared to a source sequence by generally known techniques and preferably retain the biological activity of a naturally occurring human CD16 or FcγRIII. Some variants of FcγRIII are known to those of skill in the art.

The term “CD20” refers to a cell surface protein that is expressed on certain cells of the immune system, specifically the B lymphocyte-restricted differentiation antigen Bp35. “Naturally occurring CD20” has an amino acid sequence of a protein obtained from nature and includes naturally occurring variants such as allelic

variants, isotypes and truncated forms. More specifically, "CD20" includes human CD20 (AH003353; GenBank Accession nos. M27395, J03574). Representative amino acid sequence, cDNA sequence and a genomic sequence of human and murine CD20 are shown in Figure 23. The invention also contemplates CD20

5 variants. The variants are changed as compared to a source sequence using generally known techniques and preferably retain the biological activity of a naturally occurring human CD20. Preferably the variants are not changed at amino acid positions 170 and 172 of a naturally occurring human CD20 as these positions have been shown to be a part of the epitope recognized by several different anti-

10 human CD20 monoclonal antibodies as described in Polyak et al., Blood 99:3256 (2002).

"Disruption" of a gene occurs when a fragment of DNA locates and recombines with an endogenous homologous sequence. These sequence disruptions or modifications may include insertions, missense, frameshift, deletion, or

15 substitutions, or replacements of DNA sequence, or any combination thereof. Insertions include the insertion of entire genes, which may be of animal, plant, fungal, insect, prokaryotic, or viral origin. Disruption, for example, can alter the normal gene product by inhibiting its production partially or completely or by enhancing the normal gene product's activity. In a preferred embodiment, the

20 disruption is a null disruption that has no significant expression of the gene.

By the term "endogenous loci" is meant to include the naturally occurring genetic loci found in the host animal that is to become transgenic.

The term "heterologous" when used in conjunction with polypeptide or gene refers to a polypeptide having an amino acid sequence or a DNA encoding the

25 polypeptide that is not found in transgenic nonhuman host animal. Thus, a transgenic mouse having a human CD20 gene can be described as having a heterologous CD20 gene. The transgene can be detected using a variety of methods including PCR, Western blot, or Southern blot. The term "human endogenous promoter" refers to the promoter that is naturally associated with the polynucleotide

30 sequence that encodes the human protein that is to be introduced into the animal to form a transgenic animal.

The term "non-human animals" is intended to include any vertebrate such as mammals, birds, reptiles, and amphibians. Suitable mammals include rodents, non-human primates, sheep, dogs and cows. Suitable birds include chickens, geese, and

turkeys. Preferred non-human animals are selected from the rodent family including rat and mouse, most preferably mouse.

The term "naturally-occurring" or "naturally associated" as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

For nucleic acids, the term "substantial homology" indicates that two nucleic acids, or designated sequences thereof, when optimally aligned and compared with appropriate nucleotide insertions or deletions have at least about 80% sequence identity, more preferably about 81% sequence identity, more preferably about 82% sequence identity, more preferably about 83% sequence identity, more preferably about 84% sequence identity, more preferably about 85% sequence identity, more preferably about 86% sequence identity, more preferably about 87% sequence identity, more preferably about 88% sequence identity, more preferably about 89% sequence identity, more preferably about 90% sequence identity, more preferably about 91% sequence identity, more preferably about 92% sequence identity, more preferably about 93% sequence identity, more preferably about 94% sequence identity, more preferably about 95% sequence identity, more preferably about 96% sequence identity, more preferably about 97% sequence identity, more preferably about 98% sequence identity, and more preferably about 99% sequence identity to one another. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to the complement of the strand. Methods of aligning two sequences and identifying % identity are known to those of skill in the art. Several computer programs are available for determining % identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared.

"Transcriptional regulatory sequence" refers to polynucleotide sequences, such as initiation signals, enhancers, and promoters, which induce or control transcription of protein coding sequences with which they are operably linked. In

preferred embodiments, transcription of a recombinant transgene is under the control of a promoter sequence (or other transcriptional regulatory sequence), which controls the expression of the recombinant gene in a cell-type in which expression is intended. It will also be understood that the recombinant gene can be under the control of transcriptional regulatory sequences which are the same or which are different from those sequences, which control transcription of a naturally-occurring form of CD20 or CD16.

As used herein, the term "transgene" means a nucleic acid sequence (encoding, e.g., CD20 or CD16) that has been introduced into a cell by way of human intervention such as by way of the described methods herein. A transgene could be partly or entirely heterologous, i.e., foreign, to the transgenic animal or cell into which it is introduced. A transgene can include one or more transcriptional regulatory sequences and any other nucleic acid, such as introns, that may be necessary for optimal expression of a selected nucleic acid.

"Transgenic animal" or "Tg<sup>+</sup>" are used interchangeably and are intended to include any non-naturally occurring non-human animal in which one or more of the cells of the animal contain heterologous nucleic acid encoding human CD20 and/or preferably, human CD16, that has been introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, but rather is directed to the introduction of a recombinant DNA molecule. This molecule may be integrated within a chromosome, or it may be extrachromosomally replicating DNA. The term "Tg<sup>+</sup>" includes animals that are heterozygous and/or homozygous for either human CD20 and/or human CD16.

"CD20 associated disease" refers to diseases or disorders that have been associated with the expression of CD20 on cells, aberrant proliferation or activation of B cells, or have been treated with anti-CD20 antibodies. For example, a chimeric anti-CD20 antibody has been used to treat patients with lymphoma. Other diseases or conditions that have been treated with anti-CD20 therapy include autoimmune conditions or diseases such as rheumatoid arthritis, systemic lupus erythematosus, and ankylosing spondylitis. Other conditions include Epstein Barr virus associated disease following blood or bone marrow transplant, Kaposi Sarcoma associated



herpes virus related multicentric Castlemen disease, Hepatitis C associated Cryoglobulinemia Vasculitis, lymphoproliferative disorders with autoimmune hemolytic anemia, and ANCA associated vasculitis.

5           A.     Modes of the Invention

The present invention provides transgenic animals expressing a heterologous CD20 marker, a heterologous CD16 marker or both. In one embodiment, the human CD20 transgenic animals of the invention express CD20 on the same types of B cells. Administration of anti-human CD20 antibodies to the presently described  
10 transgenic animals results in depletion of B lymphocytes expressing human CD20. Previous attempts to prepare transgenic mice expressing human CD20 may not have achieved expression of human CD20 in the appropriate B cell compartments due to lack of incorporation of effective transcriptional control regions into the transgene. In another embodiment, transgenic mice of the invention express human CD20 and  
15 human CD16.

The present invention provides a transgenic animal that has cells that react in a way that is similar to a human subject. Administration of anti-human CD20 antibodies to the presently described transgenic animals results in depletion of B lymphocytes expressing human CD20. In one embodiment, the human CD20  
20 transgenic mice are characterized by expression of human CD20 on cells at a level sufficient for anti- human CD20 antibody bound to the expressing cells to affect killing of the cells, resulting in B cell depletion of peripheral and/or circulating B cells of at least about 75% and more preferably, 80%, 85%, 90%, 95% , 99% and even 100%. A similar response is observed in humans. These animal models can  
25 be used for screening of agents including but not limited to monoclonal antibodies against the CD20 marker. In addition, transgenic mice expressing human CD16 provide models for determining an agent's ability to induce an effector cell response (such as, for example, an NK cell mediated response) or the agent's ability to further deplete B lymphocytes expressing CD20 (including malignant B cells).  
30 Additionally, the transgenic animals may be used to test the efficacy and toxicity of anti-CD20 directed therapies.

## B. DNA constructs

Typically, a polynucleotide molecule encoding a heterologous protein of the invention is inserted into a vector, preferably a DNA vector, in order to replicate the polynucleotide molecule in a suitable host cell. It will be understood that the CD20  
5 and/or CD16 transgenes may be generated and inserted separately, or generated together in a single construct for insertion. DNA constructs may also be useful to prepare targeting vectors for knockout animals.

In order to isolate, clone and transfer the CD16 and/or CD20 locus, a yeast artificial chromosome (YAC) may be employed. The entire locus can be cloned and  
10 contained within one or a few YAC clones. If multiple YAC clones are employed and contain regions of overlapping homology, they can be recombined within yeast host strains to produce a single construct representing the entire locus. YAC arms can be additionally modified with mammalian selection cassettes by retrofitting to assist in the introduction of the constructs into embryonic stems cells or embryos by  
15 the previously outlined methods.

Due to the high stability and relatively large inserts, ease of manipulation and shotgun sequencing, bacterial artificial chromosome (BAC) libraries can provide human sequences for genes of interest. BAC libraries contain an average insert size of 100-150kb. BAC clones are capable of harboring inserts as large as 300,000 base  
20 pairs. Shizuya, et al., (1992) Proc. Natl. Acad. Sci., USA 89:8794-8797; Kim, et al., (1996) Genomics 34 213-218; Swiatek, et al., (1993) Genes and Development 7:2071-2084. Genomic BAC libraries of the human and mouse have been constructed and are commercially available (Invitrogen, Carlsbad CA). Genomic BAC libraries can also serve as a source of human and murine CD20 and/or CD16  
25 gene sequences as well as transcriptional control regions.

Nucleic acids encoding human CD20 are known to those of skill in the art. Representative cDNA (SEQ ID NO: 5), genomic (SEQ ID NO: 6) and amino acid sequences (SEQ ID NO: 4) of human and murine CD20 are shown in Figure 23. Other sequences can also be found in GenBank, such as Accession No. AH003353.

30 Nucleic acids encoding the isotypes of the  $\alpha$  chain of human CD16 are known to those of skill in the art. Representative cDNA (SEQ ID NO: 2), genomic (SEQ ID NO: 3) and amino acid sequences (SEQ ID NO: 1) of human  $\alpha$  chain subtype A are shown in Figures 22A, B and C (GenBank Accession Nos. NM000569 and Z46222). Representative amino acid (SEQ ID NO: 10) and cDNA

sequences (SEQ ID NO: 11) for human CD16  $\alpha$  chain isotype B are shown in Figure 22E. (GenBank Accession No. NM000570). A genomic sequence encoding human CD16  $\alpha$  chain isotype B is shown in Figure 22F (GenBank Accession NO. Z46223). A nucleic acid sequence encoding a human Fc receptor gamma chain is also known  
5 to those of skill in the art. Representative amino acid (SEQ ID NO: 7) (GenBank Accession No. P30273) and cDNA sequences (GenBank Accession No. M33195) (SEQ ID NO: 8) are shown in Figure 25.

The heterologous transgenes preferably comprise germline regulatory DNA sequences operably linked to the gene of interest that is to be expressed in a  
10 transgenic non-human animal. By the term "operably linked" is meant a genetic sequence operationally (i.e., functionally) linked to a nucleic acid segment, or sequences upstream (5') or downstream (3') from a given segment or sequence. Those nearby sequences often impact processing and/or expression of the nucleic acid segment or sequence in a desired cell type.

15 Preferably, these regulatory sequences are genomic in origin, and include one or more introns. For example, the transgenic construct can include regulatory regions located in the 5'-flanking regions of a gene encoding CD20 and/or CD16, operably linked to the coding sequences in a manner capable of replicating and expressing the gene in a host cell. In one embodiment, the regulatory sequences  
20 comprise the endogenous promoter sequence naturally associated with either the CD20 and/or CD16. In some embodiments, the promoters provide for tissue specific expression at a level similar to that level of expression in the animal from which the sequence is derived. If additional flanking sequences are useful in optimizing expression, such sequences can be cloned using the existing sequences as  
25 probes. Additional sequences necessary for maximizing processing or expression of the transgene can be derived from genomic sequences.

Alternatively, the promoters can be those promoters associated with the corresponding endogenous gene in the transgenic host animal. For example, if the murine genes CD20 and/or CD16 genes are disrupted by integration with the  
30 corresponding human genes, the corresponding human genes are preferably integrated so as to be operably linked to murine transcriptional control regions for the endogenous murine CD20 and/or CD16 respectively.

Preferably, the regulatory sequences provide for expression of the transgene in the appropriate cells and at a level so that expression can be detected using

standard methodologies such as detection with antibodies. In one embodiment, the regulatory sequences provide for expression of the human CD20 transgene at a level at least 40% of the expression of CD20 on human cells.

5 An expression system or construct encoding a transgene can be expressed from a construct that includes transcriptional regulatory sequences specific for the CD20 marker and/or CD16 receptor, e.g., a human endogenous promoter (see, for example, U.S. Patent No. 5,877,396, incorporated herein by reference).

10 An expression system or construct encoding a transgene as described herein can also include a 3' untranslated region downstream of the DNA sequence. Such regions can stabilize the RNA transcript of the expression system and thus increases the yield of desired protein from the expression system. Among the 3' untranslated regions useful in the constructs of this invention are sequences that provide a poly A signal. Such sequences may be derived, e.g., from the SV40 small t antigen, the CD20 and/or CD16 untranslated region or other 3' untranslated sequences well  
15 known in the art.

Optionally, the expression system or construct includes a 5' untranslated region between the promoter and the DNA sequence encoding the signal sequence. Such untranslated regions can be from the same control region from which promoter is taken or can be from a different gene, e.g., they may be derived from other  
20 synthetic, semi-synthetic or natural sources.

In addition, other promoters or other transcriptional regulatory sequences not naturally associated with the transgene may be utilized. For example, heterologous promoters may provide for enhanced levels of expression or tissue specific expression. Various promoters having different strengths may be utilized as long the  
25 promoter functions in the non-human animal or in the desired tissue type. Many promoters are known to those of skill in the art.

Expression systems can be prepared using methods known in the art. For example, an expression system can be prepared as part of a larger plasmid. Such preparation allows the cloning and selection of the correct constructions in an  
30 efficient manner as is known in the art. Expression systems can be located between convenient restriction sites on the plasmid so that they can be easily isolated from the remaining plasmid sequences for incorporation into the desired mammal. Preferably, the DNA construct encoding the human CD20 and/or CD16 comprises a

bacterial artificial chromosome including the naturally associated transcriptional regulatory sequences to provide for tissue specific expression.

The various methods employed in the preparation of the plasmids and transformation of host organisms are known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see Molecular Cloning A Laboratory Manual, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989) Chapters 16 and 17.

### 10 C. Production of Transgenic Animals

Methods for generating transgenic animals of the present invention, including knock-outs and knock-ins, are well known in the art (see, generally, Gene Targeting: A Practical Approach, Joyner, ed., Oxford University Press, Inc. (2000)). Generation of the transgenic mice may optionally involve disruption of the genetic loci of the murine marker and introduction of the gene encoding the human marker into the murine genome, preferably at the same location as the endogenous gene.

According to the invention, a transgenic mouse model is generated where the human CD20 has been introduced into the murine genome (huCD20<sup>+</sup>; alternatively referenced as huCD20Tg<sup>+</sup>). An endogenous murine CD20 polypeptide is present on murine lymphocytes, but the known anti-human CD20 monoclonal antibodies do not bind to murine B cells. The endogenous murine CD20 gene, therefore, does not have to be but can be disrupted, if desired. When the endogenous murine CD20 is not disrupted then the gene encoding human CD20 is preferably inserted at a location other than that of the gene encoding murine CD20.

25 In a preferred embodiment, the genome of the transgenic animal further comprises a sequence encoding human CD16, preferably, the  $\alpha$  chain, and more preferably the subtype A alpha chain. When mice are used, knock-out lines are preferably generated wherein the murine CD16 gene, preferably the  $\alpha$  chain, has been disrupted (mCD16<sup>-/-</sup>). Separately, a transgenic murine line is generated where the human CD16 alpha chain gene has been introduced into the genome (huCD16<sup>+</sup>; 30 alternatively referenced as huCD16Tg<sup>+</sup>). The mCD16<sup>-/-</sup> and huCD16<sup>+</sup> mouse lines are then cross bred to generate a mouse line expressing human CD16 alpha chain but not expressing endogenous CD16 (huCD16<sup>+</sup>mCD16<sup>-/-</sup>). Alternatively, the human

gene can be introduced into an ES cell derived from the mCD16<sup>-/-</sup> line or can be used to disrupt the murine CD16 gene.

### **Inactivation of Endogenous Loci**

5           In a preferred embodiment, inactivation of the endogenous loci is achieved by targeted disruption through homologous recombination in embryonic stem cells. In one embodiment, DNA is introduced into a host cell and recombines at the endogenous loci to disrupt the production of endogenous CD16. Similarly, in another embodiment, DNA is introduced into a host cell and recombines at  
10   endogenous CD20 loci to disrupt production of endogenous CD20.

          The targeting construct may be produced using standard methods known in the art (see, e.g., Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; E.N. Glover (eds.), 1985, DNA Cloning: A Practical Approach, Volumes I  
15   and II; M.J. Gait (ed.), 1984, Oligonucleotide Synthesis; B.D. Hames & S.J. Higgins (eds.), 1985, Nucleic Acid Hybridization; B.D. Hames & S.J. Higgins (eds.), 1984, Transcription and Translation; R.I. Freshney (ed.), 1986, Animal Cell Culture; Immobilized Cells and Enzymes, IRL Press, 1986; B. Perbal, 1984, A Practical  
20   Guide To Molecular Cloning; F.M. Ausubel et al., 1994, Current Protocols in Molecular Biology, John Wiley & Sons, Inc.). For example, the targeting construct may be prepared in accordance with conventional ways, where sequences may be synthesized, isolated from natural sources, manipulated, cloned, ligated, subjected to in vitro mutagenesis, primer repair, or the like. At various stages, the joined  
25   sequences may be cloned, and analyzed by restriction analysis, sequencing, or the like.

          The targeting DNA can be obtained using techniques well known in the art. For example, the targeting DNA may be produced by chemical synthesis of oligonucleotides, nick-translation of a double-stranded DNA template, polymerase chain-reaction amplification of a sequence (or ligase chain reaction amplification),  
30   purification of prokaryotic or target cloning vectors harboring a sequence of interest (e.g., a cloned cDNA or genomic DNA, synthetic DNA or from any of the aforementioned combination) such as plasmids, phagemids, YACs, cosmids, BACs, bacteriophage DNA, other viral DNA or replication intermediates, or purified restriction fragments thereof, as well as other sources of single and double-stranded

polynucleotides having a desired nucleotide sequence. Moreover, the length of homology may be selected using known methods in the art. For example, selection may be based on the sequence composition and complexity of the predetermined endogenous target DNA sequence(s).

5 In one embodiment, the targeting construct of the present invention comprises a targeting region, which comprises a first sequence (or arm) homologous to a portion or region of the CD16 gene and/or CD20 to be disrupted and a second sequence homologous to a second portion or region of the gene. The targeting construct may further comprise a positive selection marker, which is preferably  
10 positioned between the first and the second DNA sequences. The positive selection marker may be operatively linked to a promoter and a polyadenylation signal.

In another embodiment, the targeting construct may contain more than one selectable marker gene, including a negative selectable marker, such as the herpes simplex virus tk (HSV-tk) gene, which is preferably positioned outside one or both  
15 of the homologous arms of the targeting construct. The negative selectable marker may be operatively linked to a promoter and a polyadenylation signal (see, e.g., U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059 and 5,631,153).

Once an appropriate targeting construct has been prepared, the targeting construct may be introduced into an appropriate host cell using any method known  
20 in the art. Various techniques may be employed in the present invention, including, for example: pronuclear microinjection; retrovirus mediated gene transfer into germ lines; gene targeting in embryonic stem cells; electroporation of embryos; sperm-mediated gene transfer; and calcium phosphate/DNA co-precipitates, microinjection of DNA into the nucleus, bacterial protoplast fusion with intact cells, transfection,  
25 polycations, e.g., polybrene, polyornithine, etc., or the like (see, e.g., U.S. Patent No. 4,873,191; Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152; Thompson et al., 1989, Cell 56:313-321; Lo, 1983, Mol Cell. Biol. 3:1803-1814; Lavitrano et al., 1989, Cell, 57:717-723). Various techniques for transforming mammalian cells are known in the art. (see, e.g., Gordon, 1989, Intl. Rev. Cytol.,  
30 115:171-229; Keown et al., 1989, Methods in Enzymology; Keown et al., 1990, Methods and Enzymology, Vol. 185, pp. 527-537; Mansour et al., 1988, Nature, 336:348-352).

Any cell type capable of homologous recombination may be used in the practice of the present invention. Examples of such target cells include cells derived

from vertebrates including mammals such as humans, bovine species, ovine species, murine species, simian species, and ether eukaryotic organisms such as filamentous fungi, and higher multicellular organisms such as plants.

Preferred cell types include embryonic stem (ES) cells, which are typically  
5 obtained from pre-implantation embryos cultured in vitro (see, e.g., Evans, M. J. et al., 1981, *Nature* 292:154-156; Bradley, M. O. et al., 1984, *Nature* 309:255-258; Gossler et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:9065-9069; and Robertson et al., 1986, *Nature* 322:445-448). The ES cells are cultured and prepared for introduction of the targeting construct using methods well known to the skilled artisan. (see, e.g.,  
10 Robertson, E. J. ed. "Teratocarcinomas and Embryonic Stem Cells, a Practical Approach", IRL Press, Washington D.C., 1987; Bradley et al., 1986, *Current Topics in Devel. Biol.* 20:357-371; by Hogan et al., in "Manipulating the Mouse Embryo": A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor N.Y., 1986; Thomas et al., 1987, *Cell* 51:503; Koller et al., 1991, *Proc. Natl. Acad. Sci. USA*, 88:10730; Dorin et al., 1992, *Transgenic Res.* 1:101; and Veis et al.,  
15 1993, *Cell* 75:229). The ES cells that will be inserted with the targeting construct are derived from an embryo or blastocyst of the same species as the developing embryo into which they are to be introduced. ES cells are typically selected for their ability to integrate into the inner cell mass and contribute to the germ line of an  
20 individual when introduced into the mammal in an embryo at the blastocyst stage of development. Thus, any ES cell line having this capability is suitable for use in the practice of the present invention.

After the targeting construct has been introduced into cells, the cells in which  
successful gene targeting has occurred are identified. Insertion of the targeting  
25 construct into the targeted gene is typically detected by identifying cells for expression of the marker gene. In a preferred embodiment, the cells transformed with the targeting construct of the present invention are subjected to treatment with an appropriate agent that selects against cells not expressing the selectable marker. Only those cells expressing the selectable marker gene survive and/or grow under  
30 certain conditions. For example, cells that express the introduced neomycin resistance gene are resistant to the compound G418, while cells that do not express the neo gene marker are killed by G418. If the targeting construct also comprises a screening marker such as GFP, homologous recombination can be identified through



screening cell colonies under a fluorescent light. Cells that have undergone homologous recombination will have deleted the GFP gene and will not fluoresce.

Alternatively, a positive-negative selection technique may be used to select homologous recombinants. This technique involves a process in which a first drug  
5 is added to the cell population, for example, a neomycin-like drug to select for growth of transfected cells, i.e. positive selection. A second drug, such as FIAU, is subsequently added to kill cells that express the negative selection marker, i.e. negative selection. Cells that contain and express the negative selection marker are killed by a selecting agent, whereas cells that do not contain and express the  
10 negative selection marker survive. For example, cells with non-homologous insertion of the construct express HSV thymidine kinase and therefore are sensitive to the herpes drugs such as gancyclovir (GANC) or FIAU (1-(2-deoxy 2-fluoro-B-D-arabinofluranosyl)-5-iodouracil). (see, e.g., Mansour et al., Nature 336:348-352: (1988); Capecchi, Science 244:1288-1292, (1989); Capecchi, Trends in Genet. 5:70-  
15 76 (1989)). Other methods include regulated positive selection (see U.S. 20030032175A1), which requires the addition of a single selective agent.

Successful recombination may be identified by analyzing the DNA of the selected cells to confirm homologous recombination. Various techniques known in the art, such as PCR and/or Southern analysis may be used to confirm homologous  
20 recombination events.

Selected cells are then injected into a blastocyst (or other stage of development suitable for the purposes of creating a viable animal, such as, for example, a morula) of an animal (e.g., a mouse) to form chimeras (see e.g., Bradley, A. in Teratocarcinomas and Embryonic Stem Cells: A Practical Approach, E. J. Robertson, ed., IRL, Oxford, pp. 113-152 (1987)). Alternatively, selected ES cells  
25 can be allowed to aggregate with dissociated mouse embryo cells to form the aggregation chimera. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Chimeric progeny harboring the homologously recombined DNA in their germ cells can be  
30 used to breed animals in which all cells of the animal contain the homologously recombined DNA. In one embodiment, chimeric progeny mice are used to generate a mouse with a heterozygous disruption in the CD16 or CD20 gene. Heterozygous transgenic mice can then be mated. It is well known in the art that typically  $\frac{1}{4}$  of the

offspring of such matings will have a homozygous disruption in the CD16 or CD20 gene.

In addition to the above described methods of inactivation of endogenous loci, additional preferred methods of inactivation are available and may include for  
5 example, use of the tet transcription system to utilize temporal control of specific genes of interest (Proc. Natl. Acad. Sci. 91:9302-9306 (1994)) or introduction of deoxycycline transcriptional regulatory controls for tissue specific control (Proc. Natl. Acad. Sci. 93:10933-10938 (1996)).

An additionally preferred method for functional inactivation includes  
10 employment of the cre-lox deletion, site specific recombination system for targeted knock-out of genetic loci, wherein loxP sites are inserted to flank genes of interest and cre recombinase activated to delete genes (Curr. Opin. Biotechnol., 5:521-527 (1994)).

Alternatively, antisense or RNAi methods may be utilized in order to inhibit  
15 transcription of the desired loci, thus resulting in functional disruption of endogenous loci (knock-down methods). In such a situation, oligonucleotides are generated which target specific sequences of the designated locus of interest, wherein successful targeting results in inhibited production of the functional protein.

Strains of knockout mice lacking murine Fc $\gamma$ RIII receptor are commercially  
20 available from Taconic. These mice lack the murine Fc $\gamma$ RIII gamma chain, which also results in loss of expression of Fc $\gamma$ RI and Fc $\gamma$ RIII. In addition, knockout mice lacking a functional murine  $\gamma$  chain and, therefore, lacking a murine Fc $\gamma$ RIII receptor can be prepared as described in U.S. Patent No. 5,877,396 which is hereby incorporated by reference.

Mice lacking a functional murine Fc $\gamma$ RIII  $\alpha$  chain can be generated using  
25 similar methods. Briefly, in one embodiment, a targeting vector can be prepared using a cDNA sequence for murine CD16  $\alpha$  chain as shown in Figure 22G (GenBank Accession No. NM010188). The targeting vector preferably includes 5' sequences upstream from the coding sequence as well as at least 300 nucleotides of  
30 the coding sequence. This construct can be cloned into a vector encoding the neomycin resistance gene such as pMC1 neo (Stratagene). The resulting vector can then be introduced into ES cells using electroporation. The ES cells are plated on neo resistant embryonic fibroblast feeder layers and then selected in the presence of G418. Selected ES cells are injected into blastocysts. Appropriate targeting of

murine CD16  $\alpha$  chain can be determined using RT-PCR or by loss of expression of murine CD16 alpha chain on cells. PCR primers can be designed using the cDNA sequence for murine CD16  $\alpha$  chain.

Knock-out mice homozygous for the loss of endogenous CD20 can be prepared as described in WO 02/062946 . Briefly, in one embodiment, CD20-deficient mice can be generated by targeted disruption of the murine CD20 gene in embryonic stem (ES) cells using homologous recombination. A targeting vector can be generated that replaces exons encoding part of the second extracellular loop, the 4<sup>th</sup> transmembrane domain, and the large carboxyl-terminal cytoplasmic domain of murine CD20 with a neomycin resistance gene. The nucleotide sequence of murine CD20 is known to those of skill in the art. (GenBank Accession No. M62541). In one embodiment, appropriate gene targeting generates an aberrant CD20 protein truncated at amino acid position 157 and fused with an 88 amino acid protein encoded by the Neo gene sequence.

After DNA transfection, neo-resistant ES cell clones carrying the targeted allele can be determined by Southern blot analysis. Cells of one ES cell clone can be injected into blastocysts that can be transferred into foster mothers. Highly chimeric male offspring (80-100% according to coat color) can be bred with C57BL/6 (B6) females for transmitting the mutation to their progeny. Mice homozygous for disruption of the CD20 gene can be obtained at the expected Mendelian frequency by crossing heterozygous offspring.

Appropriate targeting of the CD20 can be further verified by PCR analysis of genomic DNA from homozygous offspring. Presence or absence of wild type CD20 mRNA in CD20<sup>-/-</sup> mice can be confirmed by PCR amplification of cDNA generated from splenocytes of CD20<sup>-/-</sup> mice. Absence of cell surface CD20 protein expression in CD20<sup>-/-</sup> mice can be further verified by staining B220<sup>+</sup> splenocytes with murine anti-CD20 monoclonal antibodies. Targeted mutation of the CD20 gene abrogates cell surface CD20 protein expression.

### **Introducing Transgenes into Non-Human Animal**

The transgenic non-human animals of the invention are preferably produced by introducing transgenes into the germline of the animal. Embryonic target cells at various developmental stages can be used to introduce transgenes. Different methods are used depending on the stage of development of the embryonic target

cell. The specific line(s) of any animal used to practice this invention are selected for general good health, good embryo yields, good pronuclear visibility in the embryo, and good reproductive fitness. When transgenic mice are to be produced, strains such as C57BL/6 or C57BL/6 x DBA/2 F<sub>1</sub>, or FVB lines are often used  
5 (obtained commercially from Charles River Labs, Boston, Mass., The Jackson Laboratory, Bar Harbor, ME, or Taconic Labs.). Preferred strains are those with H-2<sup>b</sup>, H-2<sup>d</sup> or H-2<sup>q</sup> haplotypes such as C57BL/6 or DBA/1. Nude mice may be utilized in order to allow for the introduction of human tumor cells. Breeding and maintenance of nude mice are more difficult because of their susceptibility to  
10 infection and disease. The line(s) used to practice this invention may themselves be transgenics, and/or may be knockouts. Preferably the same line will be used for preparation of both the initial knockout mammals and the transgenic mammals. This will make subsequent breeding and backcrossing more efficient.

Introduction of the transgene into the embryo can be accomplished by any  
15 means known in the art such as, for example, microinjection, electroporation, or lipofection. For example, the transgene can be introduced into a mammal by microinjection of the construct into the pronuclei of the fertilized mammalian egg(s) to cause one or more copies of the construct to be retained in the cells of the developing mammal(s). Following introduction of the transgene construct into the  
20 fertilized egg, the egg may be incubated in vitro for varying amounts of time, or reimplanted into the surrogate host, or both. In vitro incubation to maturity is within the scope of this invention. One common method is to incubate the embryos in vitro for about 1-7 days, depending on the species, and then reimplant them into the surrogate host.

25 Reimplantation is accomplished using standard methods. Usually, the surrogate host is anesthetized, and the embryos are inserted into the oviduct. The number of embryos implanted into a particular host will vary by species, but will usually be comparable to the number of off spring the species naturally produces.

Retroviral infection can also be used to introduce transgene into a non-  
30 human animal. The developing non-human embryo can be cultured in vitro to the blastocyst stage. During this time, the blastomeres can be targets for retroviral infection (Jaenich, R. (1976) PNAS 73:1260-1264). Efficient infection of the blastomeres is obtained by enzymatic treatment to remove the zona pellucida (Manipulating the Mouse Embryo, Hogan eds. (Cold Spring Harbor Laboratory

Press, Cold Spring Harbor, 1986). The viral vector system used to introduce the transgene is typically a replication-defective retrovirus carrying the transgene (Jahner et al. (1985) PNAS 82:6927-6931; Van der Putten et al. (1985) PNAS 82:6148-6152). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-producing cells (Van der Putten, supra; Stewart et al. (1987) EMBO J. 6:383-388). Alternatively, infection can be performed at a later stage. Virus or virus-producing cells can be injected into the blastocoele (Jahner et al. (1982) Nature 298:623-628). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of the cells which formed the transgenic non-human animal. Further, the founder may contain various retroviral insertions of the transgene at different positions in the genome which generally will segregate in the offspring. In addition, it is also possible to introduce transgenes into the germ line by intrauterine retroviral infection of the midgestation embryo (Jahner et al. (1982) supra).

A third type of target cell for transgene introduction is the embryonic stem cell. Transgenes can be efficiently introduced into the ES cells by DNA transfection or by retrovirus-mediated transduction. Such transformed ES cells can thereafter be combined with blastocysts from a non-human animal. The ES cells thereafter colonize the embryo and contribute to the germ line of the resulting chimeric animal.

In one embodiment, a cDNA or genomic clone encoding human CD20 such as shown in Figure 23 can be introduced into FVB mice fertilized eggs or ES cells using microinjection or transfection. The embryos are incubated in vitro from about 1-7 days and then implanted into a surrogate host. The ES cells are combined with blastocysts from naturally mated host animals and the blastocysts are reimplanted into surrogate mothers.

In another embodiment, a cDNA or genomic clone encoding human CD16  $\alpha$  chain subtype A such as shown in Figure 22, can be introduced into ES cells or fertilized embryos used transfection or microinjection. In alternative embodiment, a cDNA or genomic clone encoding human Fc receptor gamma chain can be also introduced along with human CD16 alpha chain into ES or fertilized embryos. The fertilized eggs are incubated in vitro for 1-7 days and then reimplanted in a surrogate host. The ES cells are combined with blastocysts from naturally mated host animals and the blastocysts are reimplanted into surrogate mothers. Expression of human

CD16 alpha chain (FcγRIII α chain) in mouse cells will occur in the presence of mouse gamma chain as the mouse and human gamma chains are similar in sequence.

In one embodiment of the invention, an endogenous CD20 or CD16 gene in a nonhuman host is functionally disrupted by homologous integration of a  
5 heterologous CD20 or CD16 alpha chain gene, such that the heterologous CD20 or CD16 gene substantially replaces the endogenous CD20 or CD16 gene, respectively, and preferably completely replaces the coding sequences of the endogenous CD20 or CD16 gene. Preferably, the heterologous CD20 or CD16 gene is linked, as a  
10 consequence of homologous integration, to regulatory sequences (e.g., an enhancer promoter) of the endogenous CD20 or CD16 gene, respectively, so that the heterologous gene is expressed under the transcriptional control of regulatory elements from the endogenous CD20 or CD16 gene locus. Nonhuman hosts which are homozygous for such replacement alleles may be produced according to methods described herein. Such homozygous nonhuman hosts generally will express a  
15 heterologous CD20 or CD16, or both, but do not express the endogenous CD20 or CD16 protein. Usually, the expression pattern of the heterologous CD20 or CD16 gene will substantially mimic the expression pattern of the endogenous CD20 or CD16 gene, respectively, in the naturally-occurring (non-transgenic) nonhuman host.

20 For example, a transgenic mouse can be generated that has human CD20 gene sequences in place of endogenous murine CD20 gene sequences and which are transcriptionally controlled by endogenous murine regulatory sequences. The human CD20 generally will be expressed similarly to the murine CD20 in naturally occurring non-transgenic mice.

25 For example, a transgenic mouse can be generated that has the human CD16 α chain sequences and which are transcriptionally controlled by endogenous murine regulatory sequences. Alternatively, the human CD16 α chain can be introduced into a mouse and can then be crossed with a mouse lacking expression of a murine CD16 α chain. Expression of human CD16 α chain is expected to occur in the  
30 presence of the murine gamma chain.

Generally, a replacement-type targeting construct is employed for homologous gene replacement. Double-crossover homologous recombination between endogenous CD20 or CD16 alpha gene sequences of the targeting construct result in targeted integration of the heterologous CD20 or CD16 gene segments.

Usually, the homology targeting regions of the transgene comprise sequences which flank the endogenous CD20 and/or CD16 alpha gene segments, so that homologous recombination results in concomitant deletion of the endogenous CD20 and/or CD16 gene segments, respectively, and homologous integration of the heterologous gene segments. Substantially an entire endogenous CD20 and/or CD16 gene may be replaced with a heterologous CD20 and/or CD16 gene by a single targeting event or by multiple targeting events (e.g., sequential replacement of individual exons). One or more selectable markers, usually in the form of positive or negative selection expression cassettes, may be positioned in the targeting construct. It is usually preferred that selectable markers are located in intron regions of the heterologous replacement region.

#### Crossing of Transgenic Mice

Transgenic mice comprising transgene human CD20 can be crossed with transgenic mice comprising transgene human CD16 and lacking the murine CD16 . Preferably, the transgenic mouse comprises the human CD16  $\alpha$  chain and lacks the murine CD16  $\alpha$  chain. A manner of preparation is to generate a series of mammals, each containing one of the desired knockout constructs or transgenes. Such mammals are bred together through a series of crosses, backcrosses and selections, to ultimately generate a single mammal containing all desired knockout constructs and/or transgenes, where the mammal is otherwise congenic (genetically identical) to the wild type except for the presence of the knockout(s) constructs and/or transgene(s).

Typically, crossing and backcrossing is accomplished by mating siblings or a parental strain with an offspring, depending on the goal of each particular step in the breeding process. In certain cases, it may be necessary to generate a large number of offspring in order to generate a single offspring that contains each of the knockout constructs and/or transgenes in the proper chromosomal location. In addition, it may be necessary to cross or backcross over several generations to ultimately obtain the desired genotype.

Once the human loci have been introduced into the host genome, either by homologous recombination or random integration, and host animals have been produced with the endogenous CD16 loci inactivated by appropriate breeding of the various transgenic or mutated animals, one can produce a host which lacks the

native capability to produce endogenous CD16  $\alpha$  chain, but has the capacity to produce human CD16  $\alpha$  chain and/or CD20.

In one embodiment, transgenic mice expressing human CD16  $\alpha$  chain are mated to murine CD16  $\alpha$  chain deficient mice, thereby reconstituting expression of a specific human CD16 thereof in a mouse deficient for that CD16 polypeptide. In another embodiment, these transgenic mice can then be bred with mice expressing human CD20 to create a line of mice expressing both human CD16 and human CD20 but not endogenous CD16.

#### 10           D.     Verification of the Presence of Transgenes

Transgenic offspring of the surrogate host may be screened for the presence and/or expression of the transgene in the desired tissue, cell or animal by any suitable method. Screening is often accomplished by Southern blot or Northern blot analysis, using a probe that is complementary to at least a portion of the transgene. Western blot analysis using an antibody against the protein encoded by the transgene may be employed as an alternative or additional method for screening for the presence of the transgene product. Typically, DNA is prepared from tail tissue and analyzed by Southern analysis or PCR for the transgene. Alternatively, the tissues or cells believed to express the transgene at the highest levels are tested for the presence and expression of the transgene using Southern analysis or PCR, although any tissues or cell types may be used for this analysis.

Alternative or additional methods for evaluating the presence of the transgene include, without limitation, suitable biochemical assays such as enzyme and/or immunological assays, histological stains for particular marker or enzyme activities, flow cytometric analysis, and the like. Analysis of the blood may also be useful to detect the presence of the transgene product in the blood, as well as to evaluate the effect of the transgene on the levels of various types of blood cells and other blood constituents. In one embodiment, expression of human CD20 or CD16 or both can be detected on immune cells from spleen, bone marrow, peripheral blood, lymph nodes, and Peyer's Patches using a detectably labeled antibody to human CD20 or CD16 or both and analyzing the labeled cells using FACS.

Examination of expression patterns in the transgenic mice of the present invention reveals a mirroring of normal human CD20 expression found in human B lineage cells. Other immunological parameters examined in terms of percentage of



cells and phenotypic characteristics, including those of T cells, B cells, NK cells, macrophages, dendritic cells and neutrophils, revealed a similarity between transgenic negative and positive littermates. Examination of the expression patterns in CD20Tg+/ CD16Tg+ shows expression of both human CD20 and CD16 markers.

5

#### **E. Uses of Transgenic Animals**

Transgenic animals of the present invention represent models of CD16 and/or CD20 expression and function in humans. Accordingly, these animals are  
10 useful in studying the mechanisms behind their function and related events, and to generate and test products (e.g., antibodies, bispecifics, multispecifics etc.) useful in treating and diagnosing CD20 associated human diseases, including cancer and autoimmune conditions.

In preferred embodiments, transgenically expressed human CD20 and/or  
15 CD16 retain similar functional properties as are exhibited in human cells. For example, B lymphocytes expressing human CD20 are recognized by anti-human CD20 antibodies and as in the human, are similarly depleted from the transgenic animal in response to administration of human CD20 antibody. As described in further detail in the Examples, B lymphocytes in the transgenic mice of the present  
20 invention are depleted in response to treatment with anti-human CD20 antibodies such as Rituxan. In one embodiment, the human CD20 transgenic mice are characterized by expression of human CD20 on cells at a level sufficient for anti-human CD20 antibody bound to the expressing cells to affect killing of the cells, resulting in B cell depletion of peripheral and/or circulating B cells of at least about  
25 75% and more preferably, 80%, 85%, 90%, 95%, 99% and even 100%.

In addition, human CD20 is found on the same type of B cells as that of human CD20. Cells expressing human CD16 are also recognized by anti-human CD16 antibodies.

The transgenic CD16 animal is preferably capable of mediating at least one  
30 Fc-receptor mediated effector cell function or response. The term "Fc-receptor mediated effector cell function" is intended to include any effector function, which is triggered by binding of immunoglobulin, e.g., IgG, to an Fc receptor on an effector cell. For example, binding of an immunoglobulin, e.g., IgG, to cells bearing transgenically expressed human CD16 can induce a variety of effector functions,

such as antibody dependent cellular cytotoxicity (ADCC), NK cell mediated response and lysozyme production.

Accordingly, in one embodiment the transgenic animals of the invention are used to test agents such as antibodies, multi- or bispecific molecules, immunoadhesins (e.g., for human safety and efficacy) for binding to target epitopes, such as a region of a human CD20, human CD16 or both. Other agents can include antigen binding fragments of antibodies with or without Fc regions, single chain antibodies, minibodies (heavy chain only antibodies), heteromultimeric immunoadhesins with one of the multimers anti-human CD20 and/or anti-human CD16 antigen binding region. Other agents may include small molecules that inhibit or result in CD20-B cell depletion that may include variants of ligands of CD20 that bind to, but do not activate CD20. For example, the effectiveness of such agents to deplete CD20 expressing cells such as malignant B cells can be determined by measuring B lymphocyte levels in the transgenic animals before and after administration of the test agent.

Accordingly, the present invention provides methods of identifying agents capable of treating a B cell lymphoma, as well as agents capable of depleting or killing B lymphocytes expressing human CD20, by administering an agent to a transgenic animal that expresses human CD20 and determining whether there is a reduction in the number of B lymphocytes. As used herein, "B cell depletion" refers to a reduction in B cell levels in an animal or human after drug or antibody treatment as compared to the level before such treatment. B cell levels are measurable using well known assays as described herein. B cell depletion can be partial or complete. Preferably, the level of B cell depletion induced by the agent is an amount that correlates with a decrease or amelioration of the symptoms of the disease or disorder. The effectiveness of the putative agent (i.e., its ability to deplete B lymphocytes expressing CD20) can be assessed by measuring baseline levels of circulating B lymphocytes in a transgenic animal expressing CD20 and comparing with levels after administration in the same animal. A comparison of efficacy for B cell depletion can be made to known therapeutic agents such as anti-human CD20 antibodies (e.g. Rituxan ) to gauge effectiveness of the putative agent for treatment for CD20 associated condition. Alternatively, baseline levels of B lymphocytes can be measured in the various tissues (e.g., spleen, bone marrow, peripheral blood, lymph nodes, Peyer's Patches) of a first transgenic animal expressing human CD20.

The putative agent can then be administered to a second transgenic animal expressing human CD20. The animal is then sacrificed and the level of B lymphocytes analyzed. A reduction in the number of B lymphocytes in a transgenic animal expressing human CD20 is an indication of the agents' effectiveness in  
5 reducing cancerous cells associated with a B cell lymphoma, and/or treating a B cell lymphoma in a human subject. Combination therapies may also be tested to determine the effectiveness of depleting the desired cell types. For example, an anti-human CD20 antibody can be combined with another agent such as Br3-Fc to cause depletion of any CD20 bearing B cells that may be resistant to killing by anti-human  
10 CD20.

B lymphocyte recovery can also be assessed by measuring cell levels over time in a series of transgenic animals. The specificity of the agent for human CD20 can be assessed by comparing the agents' effect on transgenic mice expressing human CD20 with the effect on wild type mice (which do not express a CD20  
15 marker). The effectiveness of the agent can also be compared with the effect of a placebo or control substance such as a non-specific Ab or other control agent. Preferably, the agent targets most or all cells bearing human CD20 and but does not affect cells providing immune responsiveness to antigens that stimulate a T independent immune response.

20 In addition, the animals are useful models for assessing the potential immune response in a human as the initiation of an immune response in the transgenic animal upon administration of such agents is indicative that the agents will produce the same effect in humans. An effect on the immune response in such a transgenic animal can be detected for example by a change in cytokine levels, production of an  
25 antibody or a T cell response. In addition, the transgenic animal can be engineered to contain target cells (e.g. tumor, virus) prior to administration of the multi- or bispecific molecule.

The present invention thus provides methods for identifying agents capable of inducing effector cell responses such as ADCC or NK cell mediated immune  
30 responses. In particular, the invention provides methods of identifying agents capable of inducing Fc-mediated effector cell responses, specifically FcγIII-mediated effector cell responses. A putative agents' ability to induce such responses can be assessed by, for example, analyzing cytokines levels. For example, the effectiveness of the putative agent can be assessed by measuring baseline levels

of one or more cytokines associated with FcγIII-mediated effector cell responses in a first transgenic animal expressing human CD16 and comparing with levels after administration in the animal. Alternatively, a comparison can be made between a first transgenic animal expressing human CD16 to which the agent has been administered and a second transgenic animal to which either no agent has been administered or a placebo or control substance has been administered. The specificity of the agent can be assessed by comparing the agents' effect on a transgenic animal expressing human CD16 with that on a transgenic animal having a disrupted endogenous CD16 gene (i.e., a CD16 knockout) and/or with the wild-type animal (i.e., having a functional endogenous CD16). An increase in cytokine levels associated with human FcγIII-mediated effector cell responses in a transgenic animal expressing human CD16 is an indication of the agents' effectiveness in inducing such a response in a human subject.

One aspect of the invention comprises administering a putative agent to each of the human CD20 and human CD20/CD16 transgenic animals and comparing the effectiveness of the agent, for example, for killing or depletion of human CD20 B cells. One embodiment is a method of identifying an agent capable of inducing an Fc-mediated effector cell response against B lymphocytes expressing human CD20 comprising administering an agent to the CD20 transgenic animal; measuring the level of B lymphocytes expressing human CD20 in the CD20 transgenic animal; determining the percent reduction in the level of B lymphocytes; administering the agent to the CD20<sup>+</sup>/CD16<sup>+</sup> transgenic animal; measuring the level of B lymphocytes expressing human CD20 in the CD20<sup>+</sup>/CD16<sup>+</sup> transgenic animal; and determining the percent reduction in the level of B lymphocytes in the CD20<sup>+</sup>/CD16<sup>+</sup> animal; wherein if the percent reduction of B lymphocytes determined in the CD20<sup>+</sup>/CD16<sup>+</sup> animal is greater than the percent reduction determined in the CD20 animal, the agent is identified as capable of inducing an Fc-mediated effector cell response against B lymphocytes expressing human CD20.

To detect antibody binding following administration of an anti-CD16 and/or anti-CD20 antibody, or bi- or multispecific molecule to a transgenic animal, any suitable assay can be used. For example, a sample of the animal's blood can be taken and assayed for the presence of anti-CD antibody-CD complexes using screening assays known in the art, such as an enzyme-linked immunosorbent assay (ELISA), a radioimmunoassay (RIA), or a Western Blot Assay. Each of these

assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g., an antibody) specific for the complex of interest. Accordingly, in the present invention, these assays are used to detect CD-antibody complexes formed between immunoglobulins (e.g., IgG, IgA etc.) contained in the animal's blood serum and human CD markers contained on the surface of particular cells in the animal.

The CD marker-antibody complexes can be detected using e.g., an enzyme-linked antibody or antibody fragment that recognizes and specifically binds to the antibody-CD marker complexes. Alternatively, the complexes can be detected using any of a variety of other immunoassays. For example, the antibody can be radioactively labeled and used in a radioimmunoassay (RIA). The radioactive isotope can be detected by such means as the use of a gamma counter or a scintillation counter or by autoradiography.

To detect an immune response following administration of an antibody, or bi- or multispecific molecule or other agent to a transgenic animal, any suitable procedure for measuring a change in the concentration of e.g., a cytokine, antibody or T cell population in the plasma or serum of the animal can be used. For example, a change in a cytokine concentration in vivo can be detected via a variety of immunoassays, such as enzyme immunoassay (EIA), radioimmunoassay (RIA) or ELISPOT assay. Exemplary cytokines that can be assayed include: granulocyte/macrophage colony stimulating factor (GM-CSF), granulocyte colony-stimulating factor (G-CSF), macrophage colony-stimulating factor (M-CSF), interleukins 1-12 (IL-1 to IL-12).

For example, plasma can be obtained from a transgenic animal to which an antibody, bispecific, or multi specific molecule or other agent has been administered. The concentration of a cytokine can be measured using an EIA by detecting the interaction of the cytokine with an antibody, which is in turn conjugated to an enzyme. The activity of the enzyme is detected by the reaction with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means (Voller, "The Enzyme Linked Immunosorbent Assay (ELISA)", Diagnostic Horizons 2:1-7, 1978, Microbiological Associates Quarterly Publication, Walkersville, MD; Voller, et al., J. Clin. Pathol. 31:507-520 (1978); Butler, Meth. Enzymol. 73:482-523 (1981); Maggio, (ed.)

Enzyme Immunoassay, CRC Press, Boca Raton, FL, 1980; Ishikawa, et al., (eds.) Enzyme Immunoassay, Kigaku Shoin, Tokyo, 1981). Enzymes which can be used to detectably label the antibody include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, 5 alpha-glycerophosphate, dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colorimetric methods which employ a chromogenic substrate for the enzyme. Detection may also 10 be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards or using RIA.

It is also possible to label the anti-cytokine antibody or anti CD20 agent with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wave length, its presence can then be detected. Among the most 15 commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine. The antibody can also be detectably labeled using fluorescence emitting metals such as  $^{152}\text{Eu}$ , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as 20 diethylenetriaminepentaacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA). The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling 25 compounds are luminol, isoluminol, thermotropic acridinium ester, imidazole, acridinium salt and oxalate ester. Likewise, a bioluminescent compound may be used to label the antibody. Bioluminescence is a type of chemiluminescence found in biological systems in, which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined 30 by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

A non-human transgenic animal of the present invention can further provide an indication of the safety of a particular agent for administration to a human. For example, a humanized antibody or other agent can be administered to the transgenic

animal and any toxic or adverse effects as a result of the administration of the agent to the animal can be monitored as an indication of the safety and tolerability of the humanized antibody or agent for in vivo human use. Adverse events that may occur on a short term basis include headache, infection, fever, chills, pain, nausea, 5 asthenia, pharyngitis, diarrhea, rhinitis, infusion reactions, and myalgia. Short term adverse events are measured in days post treatment. Long term adverse effects include cytotoxicity of certain cell types, bleeding events due to thrombocytopenia, release of mediators due to inflammatory and/or allergic reactions, inhibition of the immune system and/or development of an anti-therapeutic agent antibody, end organ 10 toxicity, and increased incidence of infection or malignancy. Long term adverse events are measured in months post treatment.

Another aspect of the invention involves a method for determining efficacy of an anti-CD20 agent. Efficacy can be determined by administering a range of doses of the agent to set of transgenic animals having human CD20 and/or human 15 CD16 alpha chain, determining at least one dose that results in a decrease in cells bearing human CD20.

The transgenic animals of the present invention, including cells, tissues, or other materials derived therefrom, can be utilized as models for diseases, especially diseases associated or mediated by CD20 bearing cells. Animals of any species, 20 including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate disease animal models. These systems may be used in a variety of applications. Such assays may be utilized as part of screening strategies designed to identify agents, such as compounds that are capable of ameliorating disease 25 symptoms. Thus, the animal- and cell-based models may be used to identify drugs, pharmaceuticals, therapies and interventions that may be effective in treating disease.

Cell-based systems may be used to identify compounds that may act to ameliorate disease symptoms. For example, such cell systems may be exposed to a 30 compound suspected of exhibiting an ability to ameliorate disease symptoms, at a sufficient concentration and for a time sufficient to elicit such an amelioration of disease symptoms in the exposed cells. After exposure, the cells are examined to determine whether one or more of the disease cellular phenotypes has been altered to resemble a more normal or more wild-type, non-disease phenotype.

Other uses will be readily apparent to one of skill in the art.

The following non-limiting examples are illustrative of the present invention.

All documents cited herein are hereby expressly incorporated by reference.

5

## EXAMPLES

### Example 1

This example describes generation of human CD20 BAC transgenic (Tg+) mice and a study of the effects of anti-human CD20 antibody treatment in the hCD20+ mice.

10 Human CD20 transgenic mice were generated using human CD20 CITB Human BAC-D-Clone No. 117H19 from Invitrogen. (Invitrogen, Carlsbad, CA) DNA encoding human CD20 was isolated from human lymphocytes and was sent to Invitrogen. Invitrogen tested the DNA against the filters with the clones from the human BAC library and identified clone 117H19. Previous attempts to generate  
15 transgenic mice expressing human CD20 were not successful, possibly due in part to the failure to include sufficient transcriptional control regions in the transgene construct. Transgenic mice were generated by micro injecting a human CD20 BAC construct prepared with clone 117H19 into a fertilized egg of FVB inbred strain of mice. The fertilized eggs were incubated for 1-7 days and then were implanted into  
20 surrogate mice. Mice were screened based on the FACS analysis of human CD20 expression. As can be seen from the FACS plots in Figure 1, mice heterozygous (Tg+/-) and homozygous (Tg+/+) for the transgene express human CD20 on their B220+ B cells. The murine CD20 gene was not intentionally disrupted.

Figure 2 provides a schematic diagram of expression of various cell surface  
25 markers (CD43, IgM, IgD) during B cell differentiation and maturation. In the Tg+ mice, hCD20 is expressed on pre-B, immature B cells and mature B cells. Human CD20 is found on the same cell types as that of humans and is expressed on these cell types at a comparable to slightly lower level as on human B cells.

The Tg+ mice were screened for human CD20 expression in the B cells of  
30 the bone marrow, spleen, mesenteric LN and Peyer's patches using anti-human CD20 antibodies conjugated to FITC (BD Pharmingen) (results are shown in Figures 3-6). Gating the cells on B220 and CD43, CD21, or CD38 allows delineation into the various populations of B cells from the different tissues. For gating, cells were stained with anti-B220 Ab conjugated to PerCP (BD Biosciences)



and with anti-CD43 Ab, anti-CD21 Ab or anti-CD38 Ab conjugated to PE (fluorescence, Becton Dickinson).

The level of expression of human CD20 on transgenic mice peripheral blood cells was compared to that of human CD20 on human peripheral blood cells using FACS analysis and calculating mean fluorescent intensity. Peripheral blood cells were obtained from human donors and from a hCD20Tg<sup>+</sup> mice and stained with labeled anti-human CD20 antibody (mH27). The cells were analyzed by FACS and were gated on human CD19<sup>+</sup> and B220<sup>+</sup> populations. Figure 28 shows a representative comparison of the expression level of human CD20 expression on peripheral blood cells from a human CD20 transgenic mouse as compared to expression of CD20 on human peripheral blood cells. The numbers on the graph represent mean fluorescence intensity. The results show that the human CD20 on transgenic cells was expressed at a level about 40% of human CD20 on human cells.

These results show that B cells obtained from many different tissues in the transgenic mice express human CD20 marker. The human CD20 marker is found predominantly on mature B cells but also can be found on pre-B and immature B cells similar to the profile observed in humans.

The transgenic mice were then treated with anti-human CD20 mAb m2H7 in order to determine if the antibody treatment would result in B cell depletion. The antibody m2H7 can be obtained from BD PharMingen (San Diego, CA), eBioscience, or Calbiochem. The anti-human CD20 activity of m2H7 was compared to that of Rituxan in vitro assays and had comparable activity. A humanized antibody, such as Rituxan, could also be utilized in the cell killing studies because cell killing in vivo occurs over a short enough period of time that an immune response to the humanized antibody is not a concern.

The antibody m2H7 was administered to the transgenic mice as outlined in the schematic in Figure 7 at a dose of 1 mg total which is equivalent to 3.5 mg for a 70 kg person. FACS analyses were done on peripheral blood, spleen, lymph node, bone marrow, and Peyer's Patches at the days indicated by the arrows. Serum levels of anti-CD20 mAb were monitored.

Treatment of Tg<sup>+</sup> mice with anti-CD20 mAb (m2H7) alone results in depletion of B cells in peripheral blood, mature peripheral lymph node B cells, T2 and follicular B cells in the spleen (see Figures 8-11). However, it was also observed that certain B cell subsets are resistant to killing by anti-CD20 antibody

despite very high, likely saturating levels, of antibody on the cell surface. These resistant B cells are the marginal zone B cells in the spleen (Figure 10), and the germinal center B cells in both the Peyer's patches (Fig. 12) and spleen (Figure 14). In Figure 14, mice were injected with a first dose of anti-CD20 mAb at 100 ug on day 1, followed by a second, 100 ug dose on day 3 (it is likely that a single dose at 50 ug was sufficient to saturate the B cells). T2/follicular B cells were depleted but the germinal center B cells from the Peyer's patches were shown to be bound with anti-CD20 mAb but were resistant to killing.

The recovery of B cells following anti-human CD20 antibody treatment was studied. Antibody was administered to the mice at day 1. Figure 13 shows that at day 6 post antibody treatment, B cells in the peripheral blood were not detectable. At week 6, upon clearance of the antibody, hCD20+ cells begin to be detected and by week 14, B cells appeared to have recovered to normal levels. Recovery results from precursor B cells which do not express CD20, which then subsequently develop into mature B cells with human CD20+.

Figure 14 shows FACS plots demonstrating resistance of splenic germinal center B cells to short-term (single injection) anti-CD20 mAb treatment. Mice were unimmunized or immunized with sheep red blood cells (SRBC) by intraperitoneal injection at day 1 to induce germinal centers in the spleen. The germinal centers appear by day 7. At day 8, one group of mice was treated with m2H7. The control set of mice was treated with mIgG2a isotype control antibody. Spleen cells from the mice were analyzed at day 12. PNA (peanut agglutinin), which stains for germinal center, was utilized. No detectable germinal center cells were seen in the spleens of mice not immunized with SRBC whereas the spleens of immunized mice show 0.3% PNA staining cells. While T2/Follicular B cells are depleted with anti-CD20 antibody treatment, marginal center B cells in the spleen are resistant to the antibody.

Next, it was determined whether upon B cell depletion, the mice were able to develop a T independent immune response. Mice were treated with m2H7 or isotype control antibody mIgG2a at day 0. At days 3-7, B cell depletion has occurred. At day 7, the mice were injected i.v. with *Streptococcus Pneumoniae* IV to induce a response to the polysaccharide. A T cell independent response was mounted on day 11. The results shown in Figure 15 demonstrated that treatment with anti-human CD20 did not affect the B cell response from the marginal zone and

B 1 cells, i.e., the non-depleted MZ and B1 B cells confer protection to T-independent antigens. This data demonstrates that some aspects of humoral immunity, specifically T-independent B cell responses (in this case), are preserved despite treatment with anti-CD20 mAb.

5 In summary, the human CD20 transgenic animals expressed human CD20 on mature, pre-B and immature B cells in blood, bone marrow, spleen, lymph nodes and Peyer's patches. Human CD20 was expressed on transgenic cells at a level about 40% of that of CD20 expressed on human cells. Treatment of the mice with anti-human CD20 antibodies resulted in significant depletion of B cells within 3-4 days  
10 of treatment except for the marginal zone B cells in the spleen (Figure 10), and the germinal center B cells in both the Peyer's patches (Fig. 12) and spleen (Figure 14). Not to be bound by any theory, B cell death appears to be mediated by ADCC, complement dependent cytotoxicity (CDC) or apoptosis or a combination of the three. Responses to a T independent antigen were observed in anti-human CD20  
15 treated mice which is consistent with the resistance of splenic marginal zone B cells to depletion by the anti-human CD20 antibody. The B cells that are resistant to killing with anti-human CD20 antibodies provide for retention of T-independent immune response and/or an indication that combination therapy may be required to deplete all of the B cells, if desired. Recovery of B cells expressing human CD20  
20 was observed by 14 weeks after treatment with anti-human CD20 antibodies, most likely due to the maturation of precursor B cells. These results are similar to those seen in humans treated with Rituxan.

### Example 2

25 This example demonstrates the synergy between anti-CD20 mAb and BR3 antagonist treatments for B cell modulation/depletion. BR3-Fc is an immunoadhesin having the extracellular domain of human BR3 fused to constant domain of an immunoglobulin sequence in this case, human IgG1.

Human CD20 transgenic mice expressing (designated as hCD20<sup>+</sup> mice) were  
30 treated with intraperitoneal injections of anti-CD20 mAb (single injection of 100 micrograms on day 9), BR3-Fc (100 micrograms every other day from days 1 through 12), or the combination of anti-CD20 mAb and BR3-Fc. Each group consisted of 4 mice. Two days following the last injection, the mice were sacrificed

and analyzed for hCD20<sup>+</sup> B cells. FACS analysis of spleen, blood, lymph node and Peyer's Patches were analyzed for B cell markers (CD21<sup>+</sup>CD23<sup>+</sup>).

5 The results indicate that anti-CD20 mAb therapy depleted >99% of the mature circulating B cells in the blood and lymph nodes and BR3-Fc treatment decreased mature circulating B cells in the blood and lymph nodes (Figure 16). Anti-CD20 mAb therapy depleted T2 and follicular B cells, but not marginal zone B cells in the spleen, whereas BR3-Fc treatment decreased T2/follicular and marginal zone B cells in the spleen.

10 The combination of anti-CD20 mAb and BR3-Fc synergized to deplete all populations of B cells in the spleen. Anti-CD20 mAb depletion spared most of the marginal zone and some follicular/T2 splenic B cells, while BR3-Fc depletion occurred predominantly in the marginal zone and some follicular/T2 B cells (Figure 16). Thus, the combination of both reagents completely depletes B lineage cells in the spleen. Neither BR3-Fc nor 2H7 nor the combination of the two had an effect on  
15 germinal center B cells in Peyer's Patches (Figure 17). Plasma cells were not significantly affected following anti-CD20 mAb treatment (Figure 18), indicating that some aspects of immune responsiveness were maintained in mice treated with anti-CD20 antibody.

20 These results show that the combination therapy was effective to deplete most of the B cells. Similar to humans, some B cells in transgenic mice are resistant to killing with anti-CD20 antibodies. The combination therapy provide for depletion of the B cells in the spleen resistant to anti-CD20 antibodies. This shows the transgenic mice are also useful to identify combinations of agents that may be more effective in situations that have anti-CD20 resistant cells or very aggressive tumors.

25 )

### Example 3

In this experiment, it was demonstrated that natural killer cells play a role in anti-CD20 mAb mediated B cell depletion.

Hybridoma clone, which produces PK-136 mAb (specific against mouse  
5 NK1.1), was obtained from ATCC. Four groups of human CD20 transgenic mice were injected ip with control mAb, PK-136, anti-CD20 mAb and the combination of PK-136/anti-CD20, respectively. Doses of ip were as follows:

control mAb: 200ug/ip, 3 ip/week, for 1 week

PK-136: 200ug/ip, 3 ip/week, for 1 week

10 anti-CD20 mAb: 10 ug/ip, single dose

Lymphocytes from peripheral blood, lymph nodes and spleen were analyzed 3 days after anti-CD20 mAb ip. Data is expressed as mean+/-standard error, with n=8.

The results indicated that treatment with PK-136 resulted in an  
15 approximately 80% to 90% reduction in NK cell population among the tissues examined (liver, spleen and blood) (Figure 19). In the absence of majority of NK cells, 2H7 mediated B cell depletion is less efficient(Figure 20). Thus, NK cells play a role in anti-CD20 mAb mediated B cell depletion.

### 20 Example 4

Transgenic animals expressing human CD20/ CD16 were generated and evaluated for expression of the human markers.

Human CD20 transgenic mice were generated from human CD20 BAC DNA (Invitrogen, Carlsbad, CA) as described in Example 1. Mice were screened based  
25 on the FACS analysis of human CD20 expression. Nude mice with human transgene for human CD16 alpha chain subtype A and lacking murine CD16 alpha chain were obtained from Dr. Ravetch.(Rockefeller University) These mice were then mated successively with C57B16, FVB, and 129 mice to obtain mice that were positive for human transgene encoding CD16 alpha chain and were lacking murine  
30 CD16 alpha chains in a 129/nude/FVB/B6 background. The FVB mice with human CD20 were then crossed with the 129/nude/FVB/B6 mice having human CD16 alpha chain and lacking mouse CD16 alpha chain.

Expression of human markers in the huCD20Tg<sup>+</sup>huCD16Tg<sup>+</sup>mCD16<sup>-/-</sup> mice were evaluated by isolating leukocytes from peripheral blood cells and staining them

with labeled anti-human CD20 antibodies (m2H7), anti-B220 antibodies (obtained from BD PharMingen), and anti-human CD16 antibodies (obtained from BD PharMingen). Analysis of the stained cell population was conducted using FACS. The huCD20Tg<sup>+</sup>huCD16Tg<sup>+</sup>mCD16<sup>-/-</sup> were compared with CD20Tg-/Cd16Tg- (control mice), CD20Tg+/CD16Tg-, CD20Tg-/CD16Tg+ mice. The results are shown in Figure 21.

The results show that CD20Tg+/CD16Tg- mice had cells expressing human CD20 and that these cells were B cells based on their reactivity with anti-B220 antibodies. The CD20Tg-/CD16Tg+ mice had cells that expressed human CD16 based on staining with anti-human CD16 but were not B cells because these cells were not reactive with anti-B220 antibodies. The huCD20Tg<sup>+</sup>huCD16Tg<sup>+</sup>mCD16<sup>-/-</sup> had both CD20 and CD16 positive cells that appeared to be different cell populations. The results demonstrate that the CD20Tg+/CD16Tg+ mouse was successfully generated. These results are also consistent with the view that human CD20 is found on B cells and human CD16 is mainly expressed on natural killer cells, macrophages and granulocytes.

Cells expressing the human CD16 transgene were analyzed to determine which cell types expressed the transgene. Peripheral blood cells were obtained from huCD20<sup>+</sup> huCD16<sup>+</sup> mCD16<sup>-/-</sup> transgenic mice and stained with anti-human CD16 antibodies labeled with PE (BD Pharmingen), and were gated with anti-F4/80 antibodies conjugated to APC to detect macrophages or anti-DX5<sup>+</sup> antibodies to detect natural killer cells. The results are shown in Figure 24. The results show that the cells expressing the huCD16<sup>+</sup> transgene are natural killer cells and macrophages.

Cells from mice lacking murine CD16 alpha chain were analyzed to determine whether the murine CD16 alpha chain was not present on the cells. Peripheral blood cells from mice lacking murine CD16 alpha chain and wild type mice were stained with anti-mouse CD16 antibodies and analyzed by FACS. The blood cells were gated for macrophages by staining with anti-mac-1. The results are shown in Figure 26. The results show that there was no detectable murine CD16 alpha chain on cells from mice lacking CD16 alpha chain due to knockout of that gene.

Cells from mice lacking murine CD16 alpha chain were analyzed to determine whether other Fc receptors were expressed. The cells were analyzed for the presence or absence of mouse Fc gamma RI (CD64). Peripheral blood cells from

mice lacking murine CD16 alpha chain and wild type mice were stained with anti-mouse CD64 monoclonal antibody (prepared by Genentech) and analyzed by FACS. The blood cells were gated for macrophages by staining with anti-mac-1. The results are shown in Figure 27. The shaded portion is the isotype control. The results show that there was expression of Fc gamma RI (CD64) on macrophages from mice lacking CD16 alpha chain due to knockout of that gene. The weak shift in the staining may be due to the low affinity of the antibody. These results suggest that murine CD16 alpha chain -/- knockout mice expressed other mouse Fc receptors and the mouse gamma chain homodimer.

10

What is claimed is:

1. A non-human transgenic animal whose genome comprises a first nucleotide sequence encoding human CD20 and a second nucleotide sequence encoding a subunit of a heterologous FcγIII receptor.
2. The transgenic animal of claim 1 wherein said first nucleotide sequence is operably linked to a human endogenous promoter.
3. The transgenic animal of claim 2 whose cells express human CD20.
4. The transgenic animal of claim 3 wherein human CD20 is expressed on the surface of B lymphocytes.
5. The transgenic animal of claim 2, wherein said second nucleotide sequence is operably linked to a human endogenous promoter.
6. The transgenic animal of claim 1 wherein said second nucleotide sequence encodes human CD16 alpha chain subtype A.
7. The transgenic animal of claim 6 wherein said receptor is expressed on the surface of leucocytes.
8. The transgenic animal of claim 7 wherein said receptor is expressed on the surface of a cell comprising NK cells, macrophages, neutrophils, eosinophils, basophils, mast cells or thymocyte cells or mixtures thereof.
9. The transgenic animal of claim 1 wherein the genome of said animal further comprises a disruption in an endogenous gene encoding a subunit of a receptor substantially homologous to the heterologous FcγIII receptor.
10. The transgenic animal of claim 9, wherein the endogenous gene encodes a murine CD16 alpha chain.



11. A method of identifying an agent capable of treating a B cell lymphoma said method comprising:
  - a) measuring the level of B lymphocytes expressing human CD20 in an animal of claims 1 or 9;
  - b) administering said agent to the animal of claims 1 or 9; and
  - c) measuring the level of B lymphocytes expressing human CD20 in the animal;wherein a decrease in the number of B lymphocytes expressing human CD20 in the animal after treatment with the agent identifies the agent capable of treating a B cell lymphoma.
12. An agent identified according to claim 11.
13. A method of identifying an agent capable of depleting or killing cells expressing human CD20 said method comprising:
  - a) measuring the level of B lymphocytes expressing human CD20 in an animal of claims 1 or 9;
  - b) administering said agent to the animal of claims 1 or 9; and
  - c) measuring the level of B lymphocytes expressing human CD20 in the animal;wherein a decrease in the number of B lymphocytes expressing human CD20 in the animal identifies the agent as capable of depleting or killing cells expressing CD20.
14. The method of claim 13 wherein said cells are cancer cells.
15. An agent identified according to claim 14.
16. A cell or tissue derived from the transgenic animal of claim 1 or 9.
17. The transgenic animal of claim 1 or 9 wherein said animal is a rodent.
18. The transgenic animal of claim 17 wherein said rodent is a mouse.

19. A method of identifying an agent capable of inducing an Fc-mediated effector cell response said method comprising

- a) measuring the baseline level of one or more cytokines associated with an Fc-mediated effector cell response in a transgenic animal of claim 1;
  - b) administering said agent to the transgenic animal;
  - c) measuring the level of the cytokines in the animal;
- wherein an increase in the level of cytokines after administration identifies the agent as capable of inducing an Fc-mediated effector cell response.

20. A method of identifying an agent capable of inducing an Fc-mediated effector cell response against B lymphocytes expressing human CD20, said method comprising:

- a) measuring the level of B lymphocytes expressing human CD20 in a first transgenic animal ;
  - b) administering said agent to the first transgenic animal;
  - c) measuring the level of B lymphocytes expressing human CD20 in the first transgenic animal;
  - d) determining the percent reduction in the level of B lymphocytes between step (a) and step (c);
  - e) measuring the level of B lymphocytes expressing human CD20 in a second transgenic animal of claim 1;
  - f) administering said agent to the second transgenic animal of claim 1;
  - g) measuring the level of B lymphocytes expressing human CD20 in the second transgenic animal; and
  - h) determining the percent reduction in the level of B lymphocytes between step (e) and step (g);
- wherein if the percent reduction determined in step (h) is greater than the percent reduction determined in step (d), the agent is identified as capable of inducing an Fc-mediated effector cell response against B lymphocytes expressing human CD20.

21. A method of testing safety of anti- human CD20 therapy, said method comprising:

- a) measuring the level of B lymphocytes expressing human CD20 in an animal of claims 1 or 9 ;
- b) administering said agent to the animal of claims 1 or 9; and
- c) measuring the level of B lymphocytes expressing human CD20 in the animal;  
wherein a decrease in the number of B lymphocytes expressing human CD20 in the animal identifies the agent as capable of depleting or killing cells expressing CD20;
- d) monitoring the animal for short or long term adverse effects.

22. A method of testing efficacy of anti- human CD20 therapy, said method comprising:

- a) measuring the level of B lymphocytes expressing human CD20 in a set of animals of claims 1 or 9 ;
- b) administering to each animal of the set a different dose of an agent;  
and
- c) measuring the level of B lymphocytes expressing human CD20 in the animal after each dose;and
- d) determining at least one dose of the agent that results in the most B cell depletion.

# Human CD20 Transgene expression in mouse B220<sup>+</sup> cell




Figure 1

Pro-B cells —————> Pre-B —————> Immature —————> Mature





Figure 2

# Expression of human CD20 in Tg<sup>+</sup> mouse bone marrow





Figure 3

# Expression of human CD20 in Tg<sup>+</sup> mouse splenic B cells




Figure 4

Expression of human CD20 in Tg<sup>+</sup> mesenteric LNs




Figure 5



# Expression of human CD20 in Tg<sup>+</sup> Peyer's Patches





Figure 6

Effects of anti-CD20 mAb in mice

1.0 mg anti-CD20 mAb [ $\approx$ 50 mg/kg: 3.5 gm for 70 kg man]



7/42

Day -1 1 2 3 4 7 14 21

- FACS
- Peripheral Blood
- Spleen
- LN
- Bone Marrow
- Peyer's Patch

Serum levels of anti-CD20 mAb

Figure 7

# Depletion of peripheral B cells with anti-CD20 mAbs




Figure 8

# Depletion of mature peripheral LN B cells by anti-CD20 mAb




Figure 9

Depletion of splenic T2 B cells by anti-CD20 mAbs




Figure 10

# Depletion of re-circulating mature B cells by anti-CD20 mAbs




Figure 11

# Resistance of Peyer's Patches Germinal Center B cells to anti-CD20 mAbs




Figure 12

# Depletion & Recovery of B cells following anti-CD20 mAb Rx




Figure 13



# Resistance of Splenic Germinal Center B cells to short-term anti-CD20 mAb Rx




Figure 14

# Non-depleted MZ and B1 B cells confer protection to T-Independent antigens




Figure 15




Figure 16

# In Peyer's Patches




Figure 17

Plasma cells are not depleted  
following long-term anti-CD20 mAb treatment




Figure 18

# Depletion of NK cells by PK-136 mAb




Figure 19

# NK cells play a role in 2H7-mediated B cell depletion





Figure 20

FIGURE 21

Expression of Human CD20 and CD16 Transgenes





**FIGURE 22A**

MGGGAGERLFTSSCLVGLVPLGLRISLVTCP LQCGIMWQ LLLPT  
ALLLLVSAGMRTEDLPKAVVFLEPQWYRVLEKDSVTLKCQGAYS  
PEDNSTQWFHNE SL ISSQASSYFIDAATVDDSGEYRCQTNLSTL  
SDPVQLEVHIGWLLLQAPRWVFKEEDPIHLRCHSWKNTALHKVT  
YLQNGKGRKYFHHNSDFYIPKATLKDSGSYFCRGLVGSKNVSSE  
TVNITITQGLAVSTISSFFPPGYQVSFCLVMVLLFAVDTGLYFSVKT  
NIRSSTRDWKDHKFKWRKDPQDK

**FIGURE 22B**

1 gattctgtgt gtgtcctcag atgtcagcc acagacctt gagggagtaa agggggcaga  
61 cccaccacc ttgcctccag gctcttct tcttggtcct gttctatgt ggggctcct  
121 tgccagactt cagactgaga agtcagatga agttcaaga aaaggaaatt ggtgggtgac  
181 agagatgggt ggaggggctg gggaaaggct gttacttcc tctgtctag tcggtttgt  
241 cccttaggg ctccgatat ctttggtgac ttgtccttc cagtgtggca tcatgtggca  
301 gctgtcctc ccaactgct tctacttct agttcagct ggcattcgga ctgaagatct  
361 cccaaaggct gtggtgtcc tggagcctca atggtacagg gtgctcgaga aggacagtgt  
421 gactctgaag tgccaggag cctactccc tggagacaat tccacacagt ggttcacaa  
481 tgagagcctc atctcaagcc aggcctcgag ctactcatt gacgtgcca cagtgcagca  
541 cagtggagag tacagggtcc agacaaacct ctccaccctc agtgaccggg tgcagctaga  
601 agtccatata ggctggctgt tgctccaggc cctcggttg gtgtcaagg aggaagacc  
661 tattcacctg aggtgtcaca gctggaagaa cactgctcg cataaggta catattaca  
721 gaatggcaaa ggcaggaagt atttcatca taattctgac ttctacatt caaaagccac  
781 actcaaagac agcggctct actctgcag ggggctgtt gggagtaaaa atgtgtctt  
841 agagactgtg aacatcacca tactcaagg ttggcagtg tcaaccatct catcattct  
901 tccacctggg taccaagtct cttctgctt ggtgatgga ctcttttg cagtggacac  
961 aggactatat ttctgtga agacaaacat tcgaagctca acaagagact ggaaggacca  
1021 taaatttaa tgagaaagg accctcaaga caaatgacc ccatccatg ggggtaata  
1081 gagcagtagc agcagcatct ctgaacatt ctctggatt gcaaccctat catcctcag  
1141 cctctctaca agcagcagga aacatagaac tcagagccag atccctatc caactctga  
1201 ctttcttg gtctccagt gaaggga gcccagatc ttcaagcagg gaagccccag  
1261 tgagtagctg catctctaga aattgaagt tcagagctac acaaacactt ttctgtccc  
1321 aaccgtccc tcacagcaa gcaacaatac aggcctagga tggtaatct taaacatac  
1381 aaaaattgct cgtgtataa attaccagt ttagagggga aaaaaaaca attattcta  
1441 aataaatgga taagtagaat taatggtga ggcaggacca tacagagtgt ggaactgt  
1501 ggggatctag ggaattcagt gggaccaatg aaagcatggc tgagaaatag caggtagtcc  
1561 aggatagtct aaggagggtg ttccatctg agcccagaga taagggtgtc ttctagaac  
1621 attagccgta gtgaattaa caggaaatca tgagggtgac gtagaattga gtctccagg  
1681 ggactctatc agaactggac catctccaag tatataacga tgagtctct taatgctagg  
1741 agtagaaaat ggtcctagga aggggactga ggattgcgt ggggggtgg gtgaaaaga  
1801 aagtacagaa caaacctgt gtcactgtcc caagtgcta agtgaacaga actatctag  
1861 catcagaatg agaaagcctg agaagaaaga accaaccaca agcacacagg aaggaaagc  
1921 caggagggtg aaatgcttc ttggccagg tagtaagaat tagaggtaa tgcagggact  
1981 gtaaaaccac ctttctgct tcaatatcta attctgtgt agcttgttc attgattta  
2041 taaacaaat gttgtataac caatactaaa tgtactctg agcttcgctg agttaagta

**FIGURE 22B**  
**(CONT'D)**

2101 tgaaacttc aaatcctca tcatgtcagt tccaatgagg tggggatgga gaagacaatt  
2161 gttgctatg aaagaaagct ttagctgtct ctgtttgta agctttaagc gcaacatttc  
2221 ttggttcaa taaagcattt tacaagatct tgcattgctac tcttagatag aagatgggaa  
2281 aaccatggta ataaaatag aatgataaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa  
2341 a

**FIGURE 22C**

1 aagcttccca tctgttgca gtccctfact ctctctctgt gctctctct ctcttctca  
61 tctagcccac ccttttgga gctaagaatt cctccetcca ttggagagcc acagacccaaa  
121 gaggagtcaa ataagaaaat aagacctcaa agaaggaaaa caaagtgaag gccttgcac  
181 agaagtcacg tggcagaaag ccacctggat atctgaaaag aagaaagaat tgagggat  
241 ccgcttttg cctcagagac catccttagc cctgaaggct ttgtttctgc ttaggtttc  
301 ccagataagc atccgaagtg ctacagcaag gaactttaag ttccagata ctgtctgga  
361 ttttgaagg cgtagatgag tcaattgaga aggagaactg gaatggctgc ctaggttcat  
421 ttccattgtg caatccaagg gcctgtggag aaggggctgc tgaagactc tgtgtgtggc  
481 ggggggaggg gtgggtacgt ggatggcaat gggaggatca attactcca cccaggagcc  
541 aatgaaaca cacaataaaa aaacaaaacc tgagtagtgg ttttaggtc attctggagt  
601 agaaagagca ttcatftata gcaaagggtg gcgggcacct gtgcagccc ctgcctccac  
661 tccaccctca acaagtatca ggtgccaca cgggcctgct gctgcctcc tgggctttc  
721 taagccaggt gagacctgtc ccagatgtcc acgaatccac tgggggagtg gcactatcaa  
781 gcagagtcac ctgattttct gcctgggacc tggaccattg tgagagtaac caacgtggg  
841 ttacggggga gaatctggag agaagagaag aggttaacaa ccctccact tctggccac  
901 cccctccac cttttctgtt aaggagccct ggagccccgg ctctaggct gacagaccag  
961 cccagatcca gtggcccga ggggcctgag ctaaatccgc aggacctggg taacagagg  
1021 aaggtaaaga gttctgtcc tgcctctcc ccacccccc cttttctgt atctttcag  
1081 ctttctgctg gtgactgtt ctccagggc ccatttctct accctacctg ggtttctt  
1141 aacctggaaa tctaatgac aaatcacact aaaaagtcag tagctctgt ggattacata  
1201 tccaggagc atatatgtt tgaatttga atttgaaag aaattctgcg tggagataat  
1261 attgaggcag agacactgtc agtggctga agattgaaa ggaccactt ctgtgtgac  
1321 gcagggcctc agctggagat agatgggtct gggcgaggca ggagagtac aagttctgag  
1381 gtgaaatgaa ggaagccctc agagaatgct cctccacct tgaatctcat cccagggtc  
1441 tcactgtccc attcttggtg ctgggtgat ccaaatccag gagatggggc aagcatcctg  
1501 ggatggctga gggcacactc tggcagattc tgtgtgtgc ctcatgtct cagccacaga  
1561 ctttgaggg agtaaagggg gcagaccac ccacctgccc tccaggctct tcttctctg  
1621 gtctgttct atggtggggc tcccttgcca gactcagac tgagaagtca gatgaagtt  
1681 caagaaaagg aaattggtg gtgacagaga tgggtggagg ggctggggaa aggcgttta  
1741 ctctctctg tctagctgtt ttgtccct tagggctccg gatattttg gtgactgtc  
1801 cactccagtg tggcatcatg tggcagctgc tctcccaac tgcctgcta ctctaggta  
1861 agtcagggc tccctggtg agggagaagt ttgagatgcc ttgggttcag cagagacccc  
1921 tttcaggct acgaatgaga ctcccagaa gggatgggac cctcaccac atctatagct  
1981 gtggattgag ctctaggac aagccaagat ggggctagaa atgaggagaa tgcgtgtcc  
2041 aattggggca tactcatgag tgaggccagt cactcacc cctgggtcc cagaatcact

**FIGURE 22C**  
**(CONT'D)**

2101 ctgtggaacc aaagagcttc gactagatgg tcctaggggt ctgtctctt cagtttgaca  
2161 ttccaggggt ctctctatg atttcaatt tctaccctt ctgtgggga tatgggtga  
2221 ggctcttct gtagcttgg ttagggaaat tcaacctga ccctaatt gtgagttgc  
2281 acaggagca aggggtaagg gagcagtgt gaaaatagg attgtgtg acagtggcg  
2341 aagaggcatg aacagtggag accagagagc aggtagcaag gttccacca gaaacatcct  
2401 gattctggg aaaattggg tcctggggca gaggaggga ggggagttt aaactcactc  
2461 tatgttctaa tcactctgat ctctgccct actcaatatt tgatttactc tttttctg  
2521 cagtttcagc tggcatgagg actggtgagt cagcttcag gtctggatt gacctagtg  
2581 ggcacatatg gggacaaagg ccataagata ttgggaaatg ctgttgaat gggaaaatgc  
2641 tgatgtggg ttagcagga tagttctcc aacacagcag aacttgccc tgtctctc  
2701 tggccagctt tccttaagat actgaacagg ccaaaaatgg ggccaagatg ctctaagact  
2761 gagccacca gcatgggtt gcaatgagct cattctggct ttgaggctcc ctgggaatg  
2821 cagtgtagag cctgctctc tcctgtcct caccacacat tatctggct cctcagaaga  
2881 tctcccaaag gctgtggtgt tcctggagcc tcaatggtac aggtgctcg ag

## Figure 22D

## Mouse CD16 alpha chain

1 gtagttcatc tccgaacct catcagactc tgatccagtt cttgaatgac ttggacacc  
61 cagatgttic agaatgcaca ctctggaagc caatggctac ttccaccact gacaattctg  
121 ctgctgtttg cttttgcaga caggcagagt gcagctcttc cgaaggctgt ggtgaaactg  
181 gaccccccat ggatccaggt gctcaaggaa gacatggtga cáctgatgtg cgaagggacc  
241 cacaaccctg ggaactcttc taccagtggt ttccacaacg ggagggtccat ccggagccag  
301 gtccaagcca gttacacgtt taaggccaca gtcaatgaca gtggagaata tcggtgtcaa  
361 atggagcaga cccgcctcag cgaccctgta gatctgggag tgatttctga ctggctgctg  
421 ctccagaccc ctacgcgggt gtttctggaa ggggaaacca tcacgctaag gtgccatagc  
481 tggaggaaca aactactgaa caggatctca ttctccata atgaaaaatc cgtgaggat  
541 catcactaca aaagtaattt ctctatccca aaagccaacc acagtcacag tggggactac  
601 tactgcaaag gaagtctagg aagtacacag caccagtcca agcctgtcac catcactgtc  
661 caagatccag caactacatc ctccatctct ctagtctggt accacactgc ttctcccta  
721 gtgatgtgcc tctgtttgc agtggacacg ggcctttatt tctacgtacg gagaaatctt  
781 caaaccocga gggagtactg gaggaagtcc ctgtcaatca gaaagcacca ggctcctcaa  
841 gacaagtgc accccatcca tctatggca aaacatacga tgttttggtg gcagcagcaa  
901 ctttcagcc acacagcctt ctttgaaag caacttaca gcaggccggg atgtttggtt  
961 cttaatcac aacgacttag gatcaccagt tcaaggcttg ctgggtcaca cagagagagt  
1021 gagtgaagt ctacgctgga taaccagtg agatcctggg tttaggcggc tcatcaggaa  
1081 agagaacctg ttgctaattt cacaacaag atgcctactg ccatgtggc caaaggagag  
1141 aacaaggctc tggaagtgt cctctgacct ccaccatcca ccatggcagg tgcacacaat  
1201 aaattaaaat gtcattgata ttttaaaca agagacaggg gcaggctaag ggtgatggc  
1261 atagctgta tccagtacac ataatgcctt gggtttgacc tctataata aagc

## Figure 22E

## CD16 alpha chain-B

MWQLLLPTALLLLVSAGMRTE<sup>1</sup>LPKAVVFLEPQWYSVLEKDSVT  
LKCQGAYSPEDNSTQWFHNE<sup>2</sup>SLISSQASSYFIDAATVND<sup>3</sup>SGEYR  
CQTNLSTLSDPVQLEVHIGWLL<sup>4</sup>LQAPRWVFKEEDPIHLRCHSWK  
NTALHKVTYLQNGKDRKYFHH<sup>5</sup>NSDFHIPKATLKDSGSYFCRGLV  
GSKNV<sup>6</sup>SSETVNITITQGLAVSTISS<sup>7</sup>FSPPGYQVSFCLVMVLLFAVD  
TGLYFSVKT<sup>8</sup>N

1 tctttggtga cttgtccact ccagtggtgc atcatgtggc agctgctcct cccaactgct  
61 ctgctacttc tagtttcagc tggcatgagg actgaagatc tcccaaaggc tgtggtgttc  
121 ctggagcctc aatggtacag cgtgcttgag aaggacagtg tgactctgaa gtgccaggga  
181 gcctactccc ctgaggacaa ttccacacag tggtttcaca atgagagcct catctcaagc  
241 caggcctoga gctacttcat tgacgtgcc acagtcaacg acagtggaga gtacagggtg  
301 cagacaaacc tctccaccct cagtgaccgc gtgcagctag aagtccatat cggctggctg  
361 ttgctccagg ccctcgggtg ggtgttcaag gaggaagacc ctattcacct gaggtgtcac  
421 agctggaaga acactgctct gcataaggtc acatatttac agaatggcaa agacaggaag  
481 tattttcatc ataattctga ctccacatt ccaaaagcca cactcaaaga tagcggctcc  
541 tacttctgca gggggcttgt tgggagtaaa aatgtgtctt cagagactgt gaacatcacc  
601 atcactcaag gtttggcagt gtcaaccatc tcatattct ctccacctgg gtaccaagtc  
661 tctttctgct tggatggtg actcctttt gcagtggaca caggactata ttctctgtg  
721 aagacaaaca ttgaagctc aacaagagac tggaggacc ataaactta atggagaaag  
781 gaccctcaag acaaatgacc cccatcccat gggagtaata agagcagtg cagcagcatc  
841 tctgaacatt tctctgatt tgcaaccca tcatctcag gcctctc

Figure 22F

1 aagcttccca tctgttgca gtcccttact ctctctctgt gctctctct ctcttctca  
61 tctagcccac ccttttgga gctaagaatt cctccctcca ttggagagtc acagacccaaa  
121 gaggagtc aaataagaaat aagacctcaa agaaggaaaa caaagtgaag gccttgcac  
181 agaagtcacg tggcagaaaag ccacctggat atctgaaaag aagaaagaat tgagggat  
241 ccgcttttg cctcagagac catccttagc cctgaaggct ttgttctgc ttaggtttc  
301 ccagatgagc atctgaagtg ctacagcaag gaactcaag ttccagata ctgtctgga  
361 tttgcaagg cgtagatgag tcaactgaga aggagaactg gaatggcggc ctgggttcat  
421 ttccgtgtg caatccaagg gcctgtggag aaggggctgc tgcaagactc tgtgtgtggc  
481 aggggggagg gtgggtacgt ggatggcaat gggaggatca attactcca cccaggagcc  
541 aaatgaaaca cacaataaaa aaacaaaacc tgagtagtgg ttttaggtc attctggagt  
601 agaaagagca ttcaattata gcaaagggtg gcgggcacct gtgcagccc ctgcctccac  
661 tccaccccta acaagatca ggtgccaca cgggctgct gctcctcc tggtcttc  
721 taagccaagt gagacctgc ccagatgtcc acgaatccac tgggggagtg gcactatcaa  
781 gcagagtcac ctgattttc gcctgggacc tggaccattg tgagagtaac caacatggg  
841 ttacggggga gaattctggag agaagagaag aggttaacaa cctccact tctggccac  
901 cccctccac ctttctgtt aaggagccct ggagccctgg agccctggct cctaggctga  
961 cagaccagcc cagatccagt ggcccgagg ggctgagct aaatccgag gacctggga  
1021 acacaggaa ggtaaagagt tctgtctc acccctccc accccacct tttgtgat  
1081 ctttcagcc ttcactggt gactgttct tccaggccc atttctac cctacctgg  
1141 ttttctaa cctggaaatc taatgatcaa atcacactaa aaagtcagct cctgtggat  
1201 acatatcca ggagcatata gatttgaat ttgaattt gaaagaaat ctgcgtggag  
1261 ataattga ggcagagaca ctgtagtg tcaaagattt gaaaggacaa ctttctgt  
1321 gcaggcagg cctcggtg agatagatgg gtctggacga ggcaggagag tgagaagtc  
1381 tgagtgaaa tgcaggaagc cctcagaaa tgctctccc acctgaatc tcatcccg  
1441 ggtctgtg tccattctt ggtgtgggt ggaatcaat ccaggagatg ggggcaagca  
1501 tctgggaaag ctgagggcac actctggcag attctgtgtg tgcctcaga tgcagccg  
1561 cagaccttg ggggagtaa gggggcacac ccacccacct tgcctcagg ctcttctc  
1621 cctattctg ttctatggt gggctccatt gcgagactc agattgagaa atcagatgaa  
1681 gttcaagaa aaggaaactg gcaggtgaca gagatgggtg gagggactgg ggaaaggctg  
1741 ttactccct cctgtctagt cggcttggc cctttagggc tccggatct tttgtgact  
1801 gtccactcc agtgtggcat catgtggcag ctgctctcc caactgtct gctactcta  
1861 ggtaagtcag gataccctg gttgaggag aagtttgaga tgcctgggt tcatcagaga  
1921 cccctttca ggtacgaat gagactcca caaagggatg ggaccctca ccacatctat  
1981 agctgtggat tgagctacca ggacaagcca agatggggct agaatgagg agaagctg  
2041 ttcaattgg gtcatagtca tgagtgggc cagtcactc acccctctg gtccagaat



**FIGURE 22F**  
**(CONT'D)**

2101 cactatgtgg aactgaagag ctgcgactag atggcccta gggctgtct cttcagtt  
2161 gacattccag ggtctcctc tatggtttt aattctacc cttctgtg gggatatggg  
2221 ttgaggctgt ttctgtggct tggtttaggg aaattcaacc tgtacccta attgtgagt  
2281 ttgcacaggg agcaaggggt aaggagcag tgtgaaaat agggattgt gttgacagt  
2341 gcgcaagagg catgaacagt agagaccaga gagcaggtag caaggttcc accagaaaca  
2401 tctgattct tgggaaaatt gggctcctg ggcagaggag ggcaggggag tttaaactc  
2461 actctatgt ctaatcactc tgatctctgc cccactcaa tattgatt actctttt  
2521 ctgcagttt cagctggcat gcggactgt gagtcagct catggtctg gattgacca  
2581 gtggggcaca tatggggaca atggcataa gatattggga aatgctgtt gaatgggaaa  
2641 atgctgatgt ggggtagca gggatagtc ctccaacaca gcagaactg gccctgtgct  
2701 tctctggcca gcttcccta agatactgaa caggccaaa atggggccaa gatgcttaa  
2761 gactgagcca ccaagcatgg gttgcaatg agtcattct ggcttgagg ctcctggga  
2821 atggcagtg agagcctgct cctctcctg tctcaccac acattatct ggctcctcag  
2881 aagatctccc aaaggctgtg gtgttcttg agcctcaatg gtacagcgtg cttgagaagg  
2941 acagtgtgac tctgaagtgc caggagcct actccctga ggacaattcc acacagtgt  
3001 ttcaaatga gagcctcatc tcaagccagg cctcgag

## Figure 22G

## Murine FcgammaRIII

1 gtagtcac tcctgaacct catcagactc tgatccagtt ctggaatgac ttggacacc  
61 cagatgttc agaatgcaca ctctggaagc caatggctac ttccaccact gacaattctg  
121 ctgctgttg cttttgcaga caggcagagt gcagctcttc cgaaggctgt ggtgaaactg  
181 gacccccat ggatccaggt gctcaaggaa gacatggatg cactgatgtg cgaagggacc  
241 cacaaccctg ggaactcttc taccagtggt ttccacaacg ggaggccat cggagccag  
301 gtccaagcca gttacacgtt taaggccaca gtcaatgaca gtggagaata tcggtgtcaa  
361 atggagcaga cccgcctcag cgaccctgta gatctgggag tgatttctga ctggctgctg  
421 ctccagaccc ctacgagggt gtttctggaa ggggaaacca tcacgtaag gtgccatagc  
481 tggaggaaca aactactgaa caggatctca ttctccata atgaaaaac cgtgaggtat  
541 catcactaca aaagtaattt ctctatccca aaagccaacc acagtcacag tggggactac  
601 tactgcaaag gaagtctagg aagtacacag caccagtcca agcctgtcac catcactgtc  
661 caagatccag caactacatc ctccatctct ctagtctggt accacactgc ttctcccta  
721 gtgatgtgcc tctgtttgc agtggacacg ggcctttatt tctacgtacg gagaaatctt  
781 caaaccaccga gggagtactg gaggaagtcc ctgtcaatca gaaagcacca ggctcctcaa  
841 gacaagtgc accccatcca tcctatggca aaacatacga tgttttggtg gcagcagcaa  
901 ctttcagcc acacagcctt cctttgaaag caacttaca gcaggccggg atgtttggtt  
961 ctcaatcac aacgacttag gatcaccagt tcaaggcttg ctgggtcaca cagagagagt  
1021 gagtgaagt ctgacctgga taaccagtg agatcctggg tttaggcggc tcatcaggaa  
1081 agagaacctg tgctaattt cacaacaag atgcctactg cccatgtggc caaaggagag  
1141 aacaaggctc tggaagtgt cctctgacct ccaccatcca ccatggcagg tgcacacaat  
1201 aaattaaaat gtcatgtata ttttaaca agagacaggg gcaggctaag ggtgatggc  
1261 atagctgtta tccagtacac ataatgccct gggttgacc tcctataata aagc

**FIGURE 23A**

MTTPRNSVNGTFPAEPMKGPIAMQSGPKPLFRRMSSLVGPTQSF  
FMRESKTLGAVQIMNGLFHIALGGLLMIPAGIYAPICVTWYPLWGGIMYIISGSLLA  
ATEKNSRKCLVKGKMIMNSLSLFAAISGMILSIMDILNIKISHFLKMESLNFIRAHTP  
YINIYNCEPANPSEKNPSTQYCYSIQSLFLGILSVMLIFAFFQELVIAGIVENEWKR  
TCSRPKSNIVLLSAEEKKEQTIEIKEEVVGLTETSSQPKNEEDIEIPIQEEEEETE  
TNFPEPPQDQESSPIENDSSP"

**FIGURE 23B**

1 agtgtgcttg agaaacaaac tgcacccact gaactcogca gctagcatcc aaatcagccc  
61 ttgagatttg aggccttga gactcaggag tttagagagc aaaatgacaa caccagaaa  
121 ttcaataat gggacttcc cggcagagcc aatgaaaggc cctattgcta tgcaatctgg  
181 tcaaaaacca ctctcagga ggaagtcttc actggtgggc cccacgcaa gcttctcat  
241 gagggaatct aagacttgg gggctgtcca gattatgaat gggctctcc acattgccct  
301 ggggggtctt ctgatgacc cagcaggga ctatgcacc atctgtgta ctgtgtgga  
361 ccctctctgg ggaggcata tgtatattat ttccggatca ctctggcag caacggagaa  
421 aaactccagg aagtgttgg tcaaaggaaa aatgataatg aaticattga gcctcttgc  
481 tgccattctt ggaatgatc ttcaatcat ggacatactt aatattaaaa ttccattt  
541 tttaaaatg gagagtctga atttattag agctcacaca ccatatatta acatatacaa  
601 ctgtgaacca gctaaccct ctgagaaaaa ctccccatct acccaatact gtacagcat  
661 acaatctctg ttctgggca ttgtcagt gatctgac ttgcctct tccaggaact  
721 tgtaatagct ggcatcgtg agaatgaatg gaaaagaacg tgcctcagac ccaaactaa  
781 catagttctc ctgtcagcag aagaaaaaaa agaacagact attgaaataa aagaagaagt  
841 ggttgggcta actgaaacat ctccaacc aaagaatgaa gaagacattg aaattattcc  
901 aatccaagaa gaggaagaag aagaaacaga gacgaacttt ccagaacctc ccaagatca  
961 ggaatcctca ccaatagaaa atgacagctc tcttaagtg atttctctg tttctgtt  
1021 cttttttta acattagtgt tcatagcttc caagagacat gctgacttc atttctgag  
1081 gtactctgca catacgcacc acatctctat ctggccttg catggagtga ccatagctcc  
1141 ttctcttta cattgaatgt agagaatgta gccattgtag cagctgtgt tgcacgctt  
1201 ctcttttga gcaacttct tacactgaag aaaggcagaa tgagtgttc agaattgtat  
1261 ttctactaa cctgttcctt ggataggctt tttagtatag tattttttt ttgtcattt  
1321 ctccatcagc aaccaggag actgcacctg atggaaaaga tatatgactg ctcatgaca  
1381 ttctaaact atctttttt tattccacat ttattacgtttt ggtggagtcc ctttgcac  
1441 attgtttta ggatgataaa aaaaaaaaaa aaa

**FIGURE 23C**

1 tgtaaacca aagtaattgg agcgaagccc agggtagcag aagclactga ttctctgtca  
61 cctgatgtct atcagcgatt tcatcttcag gcctggacta caccactcac cctcccagtg  
121 tgcttgagaa acaaactgca ccactgaac tccgcagcta gcatccaaat cagccctga  
181 gatttgaggc ctggagact caggtaagga atcaattgc ttctttaaa tgacttaaag  
241 gaggtgatgg ataaggtata gaatggttt gaagactgga ggtcttgat ctaattcta  
301 gagttccct agtcagactt cctaatagtt ctatgactta aggaggggtg acgatatcaa  
361 ggctgcgc ccactcactc ctctaatacag tctcctctc acaattacc ctatgcagtc  
421 aactgtgaat cattccacaa aagtagtaga ttgcagcata tatattaaat catggttct  
481 aaaccattgg gtgcgaactg gagctctacc actaacaac aatataacct tgggcaaatt  
541 actaacctct aagcctcagc ttctcatcg ttaacttatt tatgtcttac atctcagaga  
601 ggggtactgt ttaacttta cagaaggata aatcgaaac taatgctcag caaagtacaa  
661 agaacaagaa tagcaacaaa aataactatt tattccaaca tgggttctt gcatacttt  
721 atttctcaa taatattat taagaagtaa ctaaatccaa aaattattt agatcctgaa  
781 caagagagaa caaaatctct acttgatgg aacttcatt ctgtggggaa gagactgaca  
841 ataagcaatt aaataaataa ggtaatttc tacagtatc aatgccgtaa agcaattaag  
901 ataggatttt gtaaaagaca gcaaatagga gtacatgta tagattgagg gtcaaggta  
961 ggctctcta ggagctgaca ttgagctac acctgaacaa aaagacacta gccatgcaca  
1021 gaccatgagc ccagttaagt gttatagcag ccacagagat aagaattatt attatttcaa  
1081 tttaacagt gaacctgagg ccagagaaat taaagaact tgccaacat ctgagaacaa  
1141 atggaggaat cactattgaa acctgggcaa tctgactcag gaggccacag tcttatatac  
1201 tgacattaga aagccttaga gagcctttc ttttcttg agaccgagtc tcaactgtt  
1261 acccaggctg gaatgcagtg gcatgatctc agctcactgc aacctcgc tctaggttc  
1321 aagcaattct cctgccttag cctcccgagt agctgggatt acaggtgcac accaacaatgc  
1381 ctggctaatt ttgtatttg tagtagagat ggggtttgc catgttagcc agcccggtct  
1441 caaacctcg acctcagggt atctgccat ctggcctcc caaagtgtt ggattacagg  
1501 catgagccac cgtgcccgac ctgagagacc ttctgatgt gactgcaca aggtggcaga  
1561 gttagagaca gagagaggcc tggaatcgac ccctctgct tctacagata gtccttacca  
1621 tactctgcaa tgtgcctct ggccatcata atgcacaaag gcagataagc aaaaggacaa  
1681 ggacaagtcc attgaaaata cattttcaa tattaagca aaagaaaagc atccaggaat  
1741 aagaaacaaa gaggacatgc agtcatatat gcaagggtgc ctctacaaag ataaagaatg  
1801 ccccaaacc agttgtcaag atcactggca gggactcctg ggccacatg ctcttctaa  
1861 acaaccctc catctcctt ctgagaactc agcagtaggc ctgcctcag atccaaggtc  
1921 actcggaaga ggccatgtct acctcaatg acactcatgg aggaaatgt gagagaagca  
1981 ttcatagatg tgacacaagg taagactgcc aaaaatctt ttctgctct cctcatttg  
2041 ttattgttt tatttttagg agtttgaga gcaaatgac aacaccaga aattcagtaa

**FIGURE 23C**  
**(CONT'D)**

2101 atgggacttt cccggcagag ccaatgaaag gccctattgc tatgcaatct ggtccaaaac  
2161 cactcttcag gaggatgtct tcactgggtg gccccacgca aagctcttc atgagggaa  
2221 ctaagacttt gggggtaagt cagttgcctt ccatcccatg tcgtagggat tctctggctg  
2281 acagaagctg atgcgggtata ggtcacatac agaattcaat ccaatttgaa gaattgggat  
2341 ccaacctgat gtctcttta tgttaacac agtggggcaa atcaggggtg catcagagaa  
2401 gttatcactt agatcacctc tgggtgatct tatgtcacct ttgggtttg gggctgtat  
2461 atgcaggggt tccccattc ccagttccat ttgccagaat ccagggcata cctgctccct  
2521 ggaaatgcc catgtggtg aggaacaga ttgaacaag aaaaagacaa aattcttggc  
2581 acctccactg ctctcttag gcattcctca cagctccaag tcaggagcca gagcttcaa  
2641 cctgtcttt gcctgctagc agtgatgatt tcagctcatc cactgctgcc tctgtctct  
2701 cccagggctg tccagattat gaatgggctc ttccacattg ccctgggggg tctctgatc  
2761 atyccagcag ggaatctatgc acctatctgt gtgactgtgt ggtaccctct ctggggaggc  
2821 attatggta gtaaaagaat agcagccatt tgggaaatgg tgcagacaaa aatgttaaaa  
2881 ggctccacag ggatatgcca gattattct gtgtgaggg aaatatatga gtaggaaata  
2941 ttattgggtt aaagtaatta agaagacagg ttgaccaa atagatataa tcccatggtt  
3001 gagagtcagt ggtcctgtt catgtgaatt cagagaaagg ggcctgcat ggatctaca  
3061 gggactgtcc aaagcaagaa ctctcaaag tcagttctgg tggggagggt ggccttagac  
3121 atttagacta gatagcaaga tgtttggaa agcaagaggc agcaggaaca tccacttcca  
3181 tctaccctt ctgtctaca attctgttg gttactatgg tacctgtga aacctgtccc  
3241 atcacaagtc agtctcattt tgcctatcga cagagcagca ctctttgac gtttatgta  
3301 catgtttcc aaatctgtaa ccctgtctgg gtgtgattg agttctgtct cttgtgtct  
3361 actatatcc tgcacagat cccagatga ttgagtaaaa ggcatgaatt tagtgtact  
3421 gagcctgaat aaaggaggaa tatgacagct gaaaaatgaa tacaactgat aaaaatgggt  
3481 ggaatggtgt gtgaaagtgt ctgaaagtgt aggtctctt ctgaccagtt atcaatgta  
3541 aaaagtgatc tctctctcc tctatctct gtctgtccc cccctctcc atctcccca  
3601 cctctcttt ttacagtata ttattccgg atcactctg gcagcaacgg agaaaaactc  
3661 caggaagtgt ttggcaagta accatatgtc ctctttccc acatgtcaga gaagtaccta  
3721 ttttttgg ttaaaaactg agacccttaa aaagccatgg tatcacagcc tctagccct  
3781 aaaaagcaaa gacctccac aatgttattg tgattttatt tatgaaaaac ttagaagcga  
3841 gatcatctga agtatgttca tgggaacaga actaaaagca gatccatgaa aaccatacct  
3901 acagtcttaa gaacgttaaa tgctgtgtga aaataataga ccttcngaa gccctatcat  
3961 ttctcccaga tcaacttta ggaaattatc tgatcaatgt catgattgat tcaaaattct  
4021 agtaagcca tttttgtc gtaacattga acaagtcagt ttacctat gtctctgagt  
4081 ttccactgg aaaggaaggg taacagtctt tgcacctg tgacgtcaa tggagtgaag  
4141 gcagtagagt gtgtgatgt gcttcacagg actataggtat ctacactgtg gtctgtccc  
4201 taaaaccctt ggggaactca tatagatcca gaggaaactg gctgcaggcg cgagogatgg  
4261 gtgaaaagag tttagcagca agtctgctcc caaaaaattc ctccccaac actgttacta

**FIGURE 23C  
(CONT'D)**

4321 aactgtgtca cticataatc aatgagggaa tgggtggatt gagatggtc ctgtcttaa  
4381 agtgacctga cacactcagt ttggggggaa aacttttat gaacatcaaa ttattctta  
4441 gatacagcca gatttactga ctigccatgt gtaggtcata gagctaggaa tgaaatatgc  
4501 gcaacataaa ataataatca ataatcccat atcattatgg tactgttat ttatatttc  
4561 ctgttcaac ctttatcat ccctgcaagg tagaacattc acactgatat tctcttacct  
4621 atgtaccca aagacatcag ccctaattgt atttggaag atagctgact ggggctgatt  
4681 gcagcctatg tcagcaggaa tagatgtgt tactgtgtt gctctgctt ttttatttc  
4741 catttatttg atagtacaga tctagagggt tctatctgaa cttcccaac ctatactta  
4801 taataccatc ccactaaagt gtgatacaag aaacttctc actctctcc cctacctat  
4861 ttatgaaggc agataataaa ctggataata ttatcttca ctattcaac aaacattat  
4921 tgagtgccta ctaggatgtt ggcagtggca gtgaaggaaa tgcaaggata caagatatag  
4981 aatcaagggt tactcttaga attttgctt tataaaacag atggatgggt aatgagatag  
5041 ggaagactga gaaacaaca ggatagagac atgatttat ttatagtga caaagagggt  
5101 aaaaagaact gagagaactt cagtataatt agttgtagt gcttgtgag tcagggcagt  
5161 tgcatttga atccctccc agattatgt ttccaaagg aaatcaaacc caattaataa  
5221 atctgtgtct ccatttcagg tcaaaggaaa aatgataatg aattcattga gcctcttgc  
5281 tgccatttct ggaatgattc ttcaatcat ggacatact aatattaaaa ttccattt  
5341 ttaaaaaatg gagagtctga atttattag agctcacaca ccatatatta acatatacaa  
5401 ctgtgaacca gctaaccct ctgagaaaaa ctcccatct acccaatact gttacagcat  
5461 acaatctctg tcttggttaa gtgtcttg taagtgtgag atggatttc tctccaggga  
5521 ggaaggatga ctgtttatt atgagcatga actctgaatt ccagaccacc tgtgttgc  
5581 tgcttcaact gattattcat acctacttt ctatcagcaa tacacattaa ccatctgtt  
5641 tgtgcaaaa gttgtgttaa gagtagggt tataaagatg ctgtctctg tactagcagt  
5701 tctcacagct attcattact tgtctaaaga atgtatctt taatcgttca attatagtca  
5761 acaataactt actgaacacc aacaatgtc ttgtgccat tattacattt tcacctcat  
5821 tctctgttg ttttcaggg catttgtca gtgatgtga tcttgcctt ctccaggaa  
5881 ctgtaatag ctggcatgt tgagaatgaa tggaaaagaa cgtgtccag acccaaact  
5941 gtaagtagta gcctccagca ccgtggtcaa tgtctgtgc cttgaagat ttattcagac  
6001 ttgagttta ataatgact tgataaggat ataagcacct gcaaaaaaat ttggcattt  
6061 aaaggcatat aataaatgac ataagtagca taaaaaccag gaggtattg ataaatgtt  
6121 gtggagattg ttgacaaagg tgcagttgt aaaagtaaag aatggttgt ttaatttct  
6181 gttttagaac atagtctcc tgcagcaga agaaaaaaa gaacagacta ttgaaataaa  
6241 agaagaagt gtgggctaa ctgaaacatc ttccaacca aagaatgaag aagacattga  
6301 aattattcca atccaagaag aggaagaaga agaaacagag acgaacttc cagaacctcc  
6361 ccaagatcag gaatcctcac caatagaaaa tgacagctct ccttaa

FIGURE<sup>37/42</sup> 23D


Mouse CD20

1 gaattccttt tttttttt ttttaaaga ttatttatt attatatgta agtacactgt  
61 agctatcttc aagtacttga gatagaagag gccaaactgat ctacactgtg agtggcta  
121 ttggccctta agccttgag ccttgagacc ttggagaccc aggcgttga aaactcaatg  
181 agtggacctt tccagcaga gcctacaaaa ggtcccctcg ccatgcaacc tgcctcaaaa  
241 gtgaacctca aaaggacatc ttactggtg ggccccacac aaagcttctt catgaggga  
301 tcaaaggctt tgggggctgt ccaaactatg aatggcctct tcatattac cctgggggga  
361 ctgctgatga tccccacagg ggtcttcgca cccatctgtt tgagtgtatg gtaccctctc  
421 tggggaggca ttatgtacat tatttcagga tcaactcctg cagctgcagc agaaaaaacc  
481 tccaggaaga gtttggtaa agcaaaagtg ataataagct ctctaagcct ctttgcctcc  
541 atttctggaa taattcttc aatcatggac atacttaaca tgacacttct tcatTTTTA  
601 aaaatgagaa gactggagct tattcaaaact tccaagccgt atgttgatat ctacgactgt  
661 gaaccatcta attcctcaga gaaaaactcc ccatctacac agtactgtaa cagcattcag  
721 tctgtgtctt tgggcattct gtcggcgatg ctgatctctg ccttctcca gaaacttg  
781 acagctggta ttgtggagaa tgagtggaaa agaattgtga ccagatccaa atctaattg  
841 gttctgctgt cagctggaga aaaaaatgag cagacgatta aaatgaaaga agaaatcatt  
901 gagctaagtg gagtatcttc ccaaccaaag aatgaagagg aaattgaaat tattccagt  
961 caggaggaag aagaagaaga agcagaaata aatttccag cacctcccca agagcaggaa  
1021 tcttgccag tggaaaatga gatcgtcct taaactctt tctttctaa gcattattgt  
1081 ttagagagct tcaagacac atagttaccc tcatctctg tggcctcca caatctattc  
1141 tccatatttt cacagctaa cttgcatag agaagccaca tctagctctc cttcacattt  
1201 gaagaatgca gtgattataa aagattgtct ttgccttgc ttagggagtc ttacactggc  
1261 agaaacgtg aagaatccaa ttctattca cctttcctt ggatgtgtgt ctcagtagtg  
1321 gtaatgggtt ttccgattt cctccatcag cagttacagc agaaggagtc agagagttca  
1381 gggaattc



FIGURE 24

Macrophages and natural killer cells express transgenic human CD16



## Figure 25A

## Fc Receptor gamma Chain

1 mipavvllll llveqaaalg epqlcyilda ilflygivlt llycrlkiqv rkaaitsyek  
61 sdgvvtglt rnqetyetlk hekppq

## Figure 25B

1 cagaacggcc gatctccagc ccaagatgat tccagcagtg gtcttgctct tactccttt  
61 ggtgaacaa gcagcggccc tgggagagcc tcagctctgc tatacctgg atgccatcct  
121 gttctgtat ggaattgtcc tcacctcct ctactgtcga ctgaagatcc aagtgcgaaa  
181 ggcagctata accagctatg agaaatcaga tgggtttac acgggcctga gcaccaggaa  
241 ccaggagact tacgagactc tgaagcatga gaaaccacca cagtagcttt agaatagatg  
301 cggtcataat ctctttggc ttctggttct tccagccctc atggttggca tcacatatgc  
361 ctgcatgcca ttaacaccag ctggccctac cctataatg atcctgtgc ctaaattaat  
421 atacaccagt ggttctcct cctgttaa gactaatgct cagatgctgt ttacggatat  
481 ttatatcta gtctactct ctgtccac ccttctctc ttcccatc ccaactccag  
541 ctaaaatatg ggaagggaga accccaata aaactgccat ggactggact c

FIGURE 26

## FACS staining of mouse CD16




Figure 27

Expression of mouse CD64 in peripheral blood of CD16<sup>-/-</sup> mice





Figure 28

# Human CD20 expression in peripheral blood



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
22 July 2004 (22.07.2004)

PCT

(10) International Publication Number  
**WO 2004/060052 A3**

(51) International Patent Classification<sup>7</sup>: **G01N 33/00**,  
C12P 21/00, A01K 67/033, 67/027, C12N 15/00

(21) International Application Number:  
PCT/US2003/039686

(22) International Filing Date:  
11 December 2003 (11.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/434,115 16 December 2002 (16.12.2002) US  
60/476,481 5 June 2003 (05.06.2003) US

(71) Applicant (for all designated States except US): **GENEN-TECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHAN, Andrew, Chee-Yuen** [US/US]; 1201, Cloud Avenue, Menlo Park, CA 94025 (US). **GONG, Qian** [CN/US]; 755 Neptune Lane, Foster City, CA 94404 (US). **MARTIN, Flavius** [RO/US]; 730 Bounty Dr., #3004, Foster City, CA 94404 (US).

(74) Agent: **BRUESS, Steven, C.**; Merchant & Gould P.C., P.O. Box 2903, Minneapolis, MN 55402-0903 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
23 September 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRANSGENIC MICE EXPRESSING HUMAN CD20 AND/OR CD16

(57) Abstract: The present invention generally relates to non-human transgenic animals expressing human cellular markers, including CD20 and/or preferably CD16.

WO 2004/060052 A3

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/39686

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : G01N 33/00; C12P 21/00; A01K 67/033, 67/027; C12N 15/00 US CL : 800/3, 4, 13-18, 21-25 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 800/3, 4, 13-18, 21-25 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EAST, BIOSIS, EMBASE, CAPLUS, SCISEARCH, MEDLINE		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,E	SERFINI et al. Characterization of CD20-transduced T lymphocytes as an alternative suicide gene therapy approach for the treatment of graft-versus-host disease. Human Gene Therapy. January 2004, Vol. 15, pages 63-76.	1-22
A	KAWANAKA et al. Expression of Fc gamma receptor III (CD16 on monocytes during hemodialysis in patients with chronic renal failure. Nephron. January 2002 Vol. 90, No. 1, pages 64-71.	1-22
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed	
Date of the actual completion of the international search 04 June 2004 (04.06.2004)		Date of mailing of the international search report 06 AUG 2004
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703) 305-3230		Authorized officer Q. Janice Li Telephone No. 703-308-0196

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**